arXiv:1110.6914v1 [math.AG] 31 Oct 2011

Primitive normal completions of the affine plane II

Pinaki Mondal

November 1, 2011

Abstract

In this article we continue from [4] the study of normal analytic compactifications of $\mathbb{C}^2$ from the point of view of their associated pencils of jets of curve germs centered at infinity. If $\bar{X}$ is a normal analytic compactification of $\mathbb{C}^2$ which is *primitive*, i.e. $\bar{X} \setminus \mathbb{C}^2$ is irreducible curve, then we show that $\bar{X}$ is projective iff $\bar{X}$ is algebraic iff at least one of the jets in the associated pencil of jets of curve-germs can be represented by a planar curve with one place at infinity. As a result we show that there are primitive normal analytic compactifications of $\mathbb{C}^2$ which are *not* algebraic. We give explicit criteria for determining if the primitive compactification corresponding to a jet of curve germs at infinity is projective or not.

1 Introduction

Disclaimer: This is an unpolished draft of the article. A clearer exposition (with more complete reference) is in order and this submission will be updated in a few days.

We freely use the notation and language of [4], which henceforth we refer to as Part I. Let δ be a divisorial semidegree on $\mathbb{C}(x,y)$ such that $\delta(x) > 0$. Then there is a degree-wise Puiseux series $\psi_\delta(x)$ in x with *finitely* many terms and a rational number $r < \mathrm{ord}_x(\psi_\delta(x))$ such that for all $f \in \mathbb{C}(x,y)$,

$$\delta(f) = \delta(x)\deg_x(f(x, \psi_\delta(x) + \xi x^r)), \tag{1}$$

where ξ is an indeterminate. Let $\Psi_\delta \in \mathbb{C}[x, x^{-1}, y]$ be the unique irreducible element such that Ψ_δ is monic in y and $(y - \psi_\delta(x))$ divides Ψ_δ, i.e. Ψ_δ is just the product of all *conjugates* of $y - \psi_\delta(x)$. Let us denote by $\mathbb{C}\langle\langle x\rangle\rangle$ the field of degree-wise Puiseux series in x. Then for all $\Phi = \sum_{q\in\mathbb{Q}, n\in\mathbb{Z}} a_{q,n}x^q y^n \in \mathbb{C}\langle\langle x\rangle\rangle[y]$, and $\omega \in \mathbb{Q}$, let us write

$$\Phi|_{>\omega} := \sum_{\substack{q\in\mathbb{Q}, n\in\mathbb{Z} \\ \delta(x^q y^n) > \omega}} a_{q,n}x^q y^n.$$

In Part I we showed that

Theorem 1.1. *ν corresponds to a primitive normal compactification of $\mathbb{C}^2$ iff $\delta(\Psi_\delta(x,y)) > 0$.*

In this article we follow up Theorem 1.1 with the following

Theorem 1.2. *Let δ, ψ_δ, r and Ψ_δ be as in Theorem 1.1. Then the following are equivalent*

1. *δ corresponds to a primitive normal algebraic compactification of $\mathbb{C}^2$.*

2. *$\delta(\Psi_\delta(x,y)) > 0$ and $\Psi_\delta|_{>\delta(\Psi_\delta)} \in \mathbb{C}[x,y]$.*

3. *$\delta(\Psi_\delta(x,y)) > 0$ and there is a plane curve C with only one place at infinity such that the degree-wise Puiseux expansion of the branch of C at the point at infinity has the form $y = \phi(x)$ with $\phi(x)|_{>r} = \psi_\delta(x)$.*

Moreover, if any of the preceding equivalent conditions hold, then the primitive normal completion of $\mathbb{C}^2$ corresponding to δ is in fact projective.

Example 1.3 (A non-algebraic normal compactification of $\mathbb{C}^2$)**.** Note that for δ as in example **??** or **??**, Ψ_δ is a polynomial in x, y. Theorem 1.2 therefore implies that in both these cases δ corresponds to a normal *projective* compactification if $\delta(\Psi_\delta(x,y)) > 0$. On the other hand, let $\psi_\delta := x^{5/2} + x^{-1}$, $r := -2$, and δ be defined in terms of ψ_δ and r as in (1). Then

$$\begin{gathered}\Psi_\delta = (y - x^{5/2} - x^{-1})(y + x^{5/2} - x^{-1}) = y^2 - x^5 - 2x^{-1}y + x^{-2},\\ \delta(\Psi_\delta) = \deg_x \left(\xi x^{-2}(2x^{5/2} + \xi x^{-2})\right) = 1/2, \text{ and}\\ \Psi_\delta|_{>1/2} = y^2 - x^5 - 2x^{-1}y\end{gathered}$$

Since $\Psi_\delta|_{>1/2}$ is not a polynomial in (x, y) even though $\delta(\Psi_\delta) > 0$, it follows from theorems 1.1 and 1.2 that δ corresponds to a primitive normal analytic compactification of $\mathbb{C}^2$ which is *not* an algebraic variety.

Theorem 1.2 in particular implies that the primitive normal compactifications of $\mathbb{C}^2$ follow a dichotomy: they are either projective algebraic, or not algebraic at all! The proof of theorem 1.2 in part II actually gives much more - it gives an explicit 'normal form' for the embeddings of every primitive algebraic compactification $\bar{X}$ of $\mathbb{C}^2$ as 'weighted complete intersections' in weighted projective spaces, which enables us to gain more detailed information about the structure of $\bar{X}$. For example one can show that the curve at infinity is non-singular and rational [**?**, Corollary **??**].

Finally, we note that 1.2 gives an answer to a question we asked in Part I, namely about effective ways to determine if a given divisorial valuation on $\mathbb{C}[x,y]$ is negative on all non-constant polynomials in $\mathbb{C}[x,y]$ or if it corresponds to a primitive normal compactification of $\mathbb{C}^2$.

Theorem 1.4. *Let ν be a divisorial discrete valuation on $\mathbb{C}[x,y]$. Let us write $\delta := -\nu$. Then ν is negative on all non-constant polynomials iff one of the following holds:*

1. *$\delta(\Psi_\delta) > 0$, or*

2. *$\delta(\Psi_\delta) = 0$ and $\Psi_\delta|_{>\delta(\Psi_\delta)} \notin \mathbb{C}[x,y]$.*

In particular, the divisorial valuations which are negative on all non-constant polynomials in $\mathbb{C}[x,y]$ but do not correspond to primitive compactifications of $\mathbb{C}^2$ are precisely those satisfying assertion 2.

Example 1.5. Let $\psi_\delta := x^{5/2} + x^{-1}$, $r := -5/2$, and δ be defined in terms of ψ_δ and r as in (1). Then

$$\begin{aligned}\Psi_\delta &= y^2 - x^5 - 2x^{-1}y + x^{-2},\\ \frac{\delta(\Psi_\delta)}{\delta(x)} &= \deg_x\left(\xi x^{-5/2}(2x^{5/2} + \xi x^{-5/2})\right) = 0, \text{ and}\\ \Psi_\delta|_{>0} &= y^2 - x^5 - 2x^{-1}y \notin \mathbb{C}[x,y].\end{aligned}$$

Therefore Theorem 1.4 implies that δ is positive on all non-constant polynomials in (x, y), even though δ does not correspond to any primitive normal compactification of $\mathbb{C}^2$ (which follows from Theorem 1.1).

2 Divisorial Semidegrees on $\mathbb{C}[x, y]$

Definition 2.1. Let k be a field, A be a k-algebra and ν be a valuation on A such that $\nu(k^*) = 0$, where $k^* := k \setminus \{0\}$. Recall the following terminology from valuation theory:

- The *rank* $\operatorname{Rank}(\nu)$ of ν is the Krull dimension of the valuation ring R_ν of ν in the quotient field K of A.
- The *rational rank* of ν is $\operatorname{rat.rank}(\nu) := \dim_{\mathbb{Q}}(\nu(K^*) \otimes_{\mathbb{Z}} \mathbb{Q})$, where $K^* := K \setminus \{0\}$.
- The *transcendence degree* $\operatorname{tr.deg}_k(\nu)$ of ν (with respect to k) is the transcendence degree of the field $R_\nu/\mathfrak{m}_\nu$ over k, where $\mathfrak{m}_\nu$ is the maximal ideal of R_ν.
- Assume $\operatorname{tr.deg}(K/k) < \infty$. Then ν is called *divisorial* (with respect to k) iff $\operatorname{tr.deg}_k(\nu) = \operatorname{tr.deg}(K/k) - 1$.

At first we recall some preliminary results about divisorial valuations and valuations on the polynomial ring in two variables.

Theorem 2.2 ([6, Theorem VI.31])**.** *Let A and k be as in definition 2.1. If ν is a divisorial valuation on A with respect to k, then* $\operatorname{Rank}(\nu) = \operatorname{rat.rank}(\nu) = 1$. *In particular, the value group of ν is a subgroup of $\mathbb{R}$.* ☐

Lemma 2.3. *Let K/k be a field extension with finite transcendental degree and L be a finite algebraic extension of K. Then for each valuation ν on L which is divisorial with respect to k, the restriction of ν to K is also divisorial.*

Proof. Let R_ν be the valuation ring of ν in L and $\mathfrak{m}_\nu$ be the maximal ideal of R_ν. Then the valuation ring of $\nu|_K$ is $R_\nu \cap K$ and its maximal ideal is $\mathfrak{m}_\nu \cap K$. Since L is a finite algebraic extension of L, it follows that $R_\nu/\mathfrak{m}_\nu$ is a finite algebraic extension of $(R_\nu \cap K)/(\mathfrak{m}_\nu \cap K)$ [6, Section VI.6, Lemma 2, Corollary 2]. Therefore $\operatorname{tr.deg}_k(\nu) = \operatorname{tr.deg}_k(\nu|_K)$, which proves the lemma. ☐

Theorem 2.4 ([1, Proposition 4.1, Theorem 4.17])**.** *Let ν be an $\mathbb{R}$-valued valuation on $\mathbb{C}[u, v]$ with $\nu(u) = 1$ and $\nu(v) > 0$. Then ν can be extended to a valuation $\nu^* : \hat{k}[v] \to \bar{\mathbb{R}}_+$, where $\hat{k}$ is the field of Puiseux series in u and $\bar{\mathbb{R}}_+ := \mathbb{R}_{\geq 0} \cup \{\infty\}$. If ν is divisorial (with respect to $\mathbb{C}$), then there is $\hat{\psi} \in \hat{k}$ and a rational number r such that for all $\phi \in \hat{k}$, $\nu^*(v - \phi) = \min\{r, \operatorname{ord}_u(\hat{\psi} - \phi)\}$.* ☐

Let ν be a divisorial valuation on $\mathbb{C}[u,v]$. According to theorem 2.2 we may assume that the value group of ν is contained in $\mathbb{R}$. Assume that $\nu(u) = 1$. Let $\hat{\psi}, r$ be as in theorem 2.4 and let ψ be the *truncation* of $\hat{\psi}$ at r, i.e. if $\hat{\psi} = \sum_{l \geq m} a_l u^{l/p}$, then $\psi = \sum_{pr > l \geq m} a_l u^{l/p}$. In particular, ψ is a *finite* Puiseux series with $\deg_u(\psi) < r$. Let ξ be an indeterminate and $\psi^* := \psi + \xi u^r$. A direct application of Theorem 2.4 gives the following

Corollary 2.5. *For all* $f \in \hat{k}[v]$, $\nu(\phi(u,v)) = \operatorname{ord}_u(\phi(u, \psi^*(u,\xi)))$.

Proof. Theorem 2.4 implies that $\nu^*(v - \tau) = \operatorname{ord}_u(\psi^* - \tau)$ for all $\tau \in \hat{k}$. Since every element ϕ in $\hat{k}[v]$ factors as a product of an element of $\hat{k}$ and elements of the form $v - \tau$ for $\tau \in \hat{k}$, the corollary immediately follows. $\square$

Definition 2.6. Let A and k be as in definition 2.1. We call a semidegree δ on A *divisorial* if $-\delta$ is a divisorial valuation on A.

Corollary 2.7. *Let δ be a divisorial semidegree on $\mathbb{C}[x,y]$ such that $\delta(x) > 0$. Then there is a finite Puiseux series ψ in x and a rational number $r < \operatorname{ord}_x(\psi)$ such that for all $f \in \mathbb{C}[x,y]$,*

$$\delta(f) = \delta(x) \deg_x(f(x, \psi(x) + \xi x^r)). \tag{2}$$

Proof. Pick a positive integer q such that $\delta(x^q) = q\delta(x) > \delta(y)$. Let $u := 1/x$, $v := y/x^q$ and ν be the restriction of $-\delta$ to $\mathbb{C}[u,v]$. Since $\mathbb{C}(u,v) = \mathbb{C}(x,y)$, it follows that $\operatorname{tr.deg}(\nu) = \operatorname{tr.deg}(-\delta) = 1$. Therefore ν is a divisorial valuation, and corollary 2.5 implies that there is a finite Puiseux series ψ in u and a rational number $r > \deg_u(\psi)$ such that

$$\frac{\nu(\phi(u,v))}{\nu(u)} = \operatorname{ord}_u(\phi(u, \psi(u) + \xi u^r))$$

for all $\phi \in \hat{k}[v]$. It follows that, for all $f \in \mathbb{C}[x,y]$,

$$\begin{aligned}
\delta(f(x,y)) &= -\nu(f(x,y)) \\
&= -\nu(f(1/u, v/u^q)) \\
&= -\nu(u) \operatorname{ord}_u(f(1/u, (\psi(u) + \xi u^r)/u^q)) \quad (\text{since } f(1/u, v/u^q) \in \hat{k}[v]) \\
&= -\delta(x) \operatorname{ord}_u(f(1/u, (\psi(u) + \xi u^r)/u^q)) \quad (\text{since } \delta(x) = \nu(u)) \\
&= \delta(x) \deg_x(f(x, x^q \psi(1/x) + \xi x^{q-r}))
\end{aligned}$$

Since $\psi(u)$ is a finite Puiseux series in u, it follows that $\psi(1/x)$ is a finite Puiseux series in x. Moreover, $\operatorname{ord}_x(x^q \psi(1/x)) = q + \operatorname{ord}_x(\psi(1/x)) = q - \deg_u(\psi(u)) > q - r$ (since $r > \deg_u(\psi(u))$). Therefore the proof of the corollary is complete. $\square$

Remark - definition 2.8. For a Puiseux series ϕ in x, the *polydromy order* of ϕ is the smallest positive integer p such that $\phi \in \mathbb{C}(x^{1/p})$. Set $t := x^{1/p}$, ζ to be a p-th root of unity and ϕ^* to be the presentation of ϕ as a Puiseux series in t, i.e. $\phi^*(t) := \phi(t^p)$. Then the *conjugates* of ϕ are $\phi_j := \phi^*(\zeta^j t)$ for $0 \leq j \leq p-1$. Let $K(x)$ be the field of Puiseux series in x. Then $K(x)$ is algebraically closed. Moreover, if $f \in \mathbb{C}[x,y]$ and $y - \phi$ divides f in $K(x)[y]$, then for each j, $0 \leq j \leq p-1$, $y - \phi_j$ also divides f in $K(x)[y]$.

Remark 2.9. Let δ and ψ be as in corollary 2.7. Recall that. If the polydromy order p of ψ is greater than 1, then ψ is *not* unique. Indeed, each conjugate of ψ defines the same semidegree δ as ψ (with the same r as the one associated to ψ).

Remark 2.10. Let ψ be a degree-wise Puiseux series in x with coefficients being polynomials in an indeterminate ξ, i.e. $\psi \in \mathbb{C}\langle\langle x\rangle\rangle[\xi]$. Then the *$x$-invariant degree* $\deg_{\mathrm{x}}^{\mathrm{inv}}(\psi)$ of ψ is the largest exponent ω of x such that the coefficient of x^{ω} in ψ is a non-constant polynomial in ξ. If the Puiseux series associated to two semidegrees are conjugates and their x-invariant degrees are same, then the semidegrees are proportional. The difference $\deg_x(\psi) - \deg_{\mathrm{x}}^{\mathrm{inv}}(\psi)$ is called the *excess degree* of ψ and denoted as $\epsilon(\psi)$. If $\psi_1, \ldots, \psi_k \in \mathbb{C}\langle\langle x\rangle\rangle[\xi]$, then

$$\epsilon\left(\prod_{j=1}^{k} \psi_j\right) = \min_{j=1}^{k} \epsilon\left(\psi_j\right) \tag{3}$$

Example 2.11. Let δ be the iterated semidegree from example **??**. Then $\delta(y^2 - x^3) = 1$. Since $y^2 - x^3 = (y - x^{3/2})(y + x^{3/2})$, it follows that there are two choices for ψ, namely $\psi_0(x) := x^{3/2}$ and $\psi_1(x) := -x^{3/2}$. In both cases $r = -\frac{1}{2}$.

Example 2.12. It is possible for ψ to be zero. In fact, $\psi \equiv 0$ iff δ is a weighted degree in (x, y)-coordinates such that $\delta(x) > 0$ and $\delta(y) = r\delta(x)$.

Definition 2.13. A formal (possibly infinite) linear combination $\phi(x) := \sum_{j=1}^{\infty} a_j x^{q_j}$ is called a *degree-wise Puiseux series* in x if $q_1 > q_2 > \cdots$ is a *decreasing* sequence of rational numbers and $\phi(1/x)$ is a usual Puiseux series in x. We write $K^*(x)$ for the field of degree-wise Puiseux series in x. The *polydromy order* of $\phi(x)$ is defined to be that of $\phi(1/x)$. Similarly, the *conjugates* of ϕ are precisely those degree-wise Puiseux series η such that $\eta(1/x)$ are conjugates of $\phi(1/x)$. If the polydromy order of ϕ is greater than 1, then a collection $(p_1, q_1), \ldots, (p_l, q_l)$ of pairs of integers are called *degree-wise Puiseux pairs* of ϕ iff $(p_1, -q_1), \ldots, (p_l, -q_l)$ are Puiseux pairs of $\psi(1/x)$. More precisely, $(p_1, q_1), \ldots, (p_l, q_l)$ are degree-wise Puiseux pairs of ϕ if and only if

1. $p_1 > 1$, and for each i, $1 \le i \le k$, p_i is a positive integer and $\gcd(p_i, q_i) = 1$, and
2. ϕ has an expression of the following form:

$$\begin{aligned}\phi = \sum_{j=1}^{k_0} a_{0j} x^{q_{0j}} + \left(a_1 x^{\frac{q_1}{p_1}} + \sum_{j=1}^{k_1} a_{1j} x^{\frac{q_{1j}}{p_1}}\right) + \left(a_2 x^{\frac{q_1}{p_1 p_2}} + \sum_{j=1}^{k_2} a_{2j} x^{\frac{q_{2j}}{p_1 p_2}}\right) + \\ \cdots + \left(a_{l-1} x^{\frac{q_{l-1}}{p_1 p_2 \cdots p_{l-1}}} + \sum_{j=1}^{k_{l-1}} a_{l-1,j} x^{\frac{q_{l-1,j}}{p_1 p_2 \cdots p_{l-1}}}\right) + a_l x^{\frac{q_l}{p_1 p_2 \cdots p_l}} + \sum_{j=1}^{\infty} a_{lj} x^{\frac{q_{lj}}{p_1 p_2 \cdots p_l}}\end{aligned} \tag{4}$$

   such that

   (a) the exponents of x in the expression in (4) are strictly decreasing,
   (b) k_i's are non-negative integers for all i, $0 \le i \le l - 1$,
   (c) q_{ij}'s are integers for all i, j, $1 \le i \le l$, $1 \le j \le k_i$, and
   (d) a_i's are nonzero complex numbers for all i, $1 \le i \le k$

   In particular, the polydromy order of ϕ is $p_1 p_2 \cdots p_l$.

Remark 2.14. Since the field of Puiseux series in x is algebraically closed, it follows that field of degree-wise Puiseux series $K^*(x)$ in x is also algebraically closed. Moreover, for all $f \in \mathbb{C}[x,y]$, remark-definition 2.8 implies that there is a unique factorization of f in $K^*(x)[y]$ of the form

$$f = x^e \left(\prod_{j=0}^{m_1-1} (y - \phi_{1j}) \right)^{e_1} \cdots \left(\prod_{j=0}^{m_n-1} (y - \phi_{nj}) \right)^{e_n} \tag{5}$$

such that $n, e, e_1, \ldots, e_n$ are non-negative integers, $\{\phi_{i0} : 1 \leq i \leq n\}$ is a collection of mutually non-conjugate degree-wise Puiseux series, and for each i, $1 \leq i \leq n$, ϕ_{ij}'s are the conjugates of ϕ_{i0}.

Definition 2.15. Let $\phi = \sum_j a_j x^{q_j/p}$ be a degree-wise Puiseux series with polydromy order p and r be a multiple of p. Then for all $c \in \mathbb{C}$ we define

$$c \star_r \phi := \sum_j a_j c^{q_j r/p} x^{q_j/p}.$$

Remark 2.16. With ϕ and p as in definition 2.15, note that if d and e are positive integers and $c \in \mathbb{C}$, then $c \star_{pde} \phi = c^e \star_{pd} \phi = c^{de} \star_p \phi$.

Notation 2.17. If ϕ and ψ are two degree-wise Puiseux series in x and $r \in \mathbb{Q}$, we write $\phi \equiv_r \psi$ iff $\deg(\phi - \psi) \leq r$.

Remark - Notation 2.18. If ϕ is a degree-wise Puiseux series in x with polydromy order p, then we write

$$\Phi^* := \prod_{j=1}^{p} (y - \zeta^j \star_p \phi(x)),$$

where ζ is a p-th root of unity. Note that Φ is a degree-wise Puiseux series in x with coefficients being polynomials in y. If $(p_1, q_1), \ldots, (p_l, q_l)$ are Puiseux pairs of ϕ, then for each k, $1 \leq k \leq l$, we write

$$\Phi^{(k)} = \prod_{j=1}^{p_1 \ldots p_k} (y - \zeta^j \star_p \phi(x)),$$

where ζ_k is a $p_1 \ldots p_k$-th root of unity.

Let δ be a divisorial semidegree on $\mathbb{C}[x,y]$ such that $\delta(f)$ is positive for all non-constant $f \in \mathbb{C}[x,y]$. Then according to corollary 2.7, there is a finite Puiseux series ψ in x and a rational number $r < \mathrm{ord}_x(\psi)$ such that (2) holds for all $f \in \mathbb{C}[x,y]$. Changing y by $y - h(x)$ for a polynomial $h(x) \in \mathbb{C}[x]$, we may assume that ψ has the form

$$\psi = \left(a_1 x^{\frac{q_1}{p_1}} + \sum_{j=1}^{k_1} a_{1j} x^{\frac{q_{1j}}{p_1}} \right) + \cdots + \left(a_l x^{\frac{q_l}{p_1 p_2 \cdots p_l}} + \sum_{j=1}^{k_l} a_{l,j} x^{\frac{q_{l,j}}{p_1 p_2 \cdots p_l}} \right) \tag{6}$$

where $(p_1, q_1), \ldots, (p_l, q_l)$ are degree-wise Puiseux pairs of ψ. Note that

$$q_1 \text{ is positive, since } \delta(y) \text{ is positive.}$$

Let δ_0 be the weighted degree on G corresponding to weights 1 for x and q_1/p_1 for y. Set $p_{l+1} := 1$, $p := p_1 p_2 \cdots p_{l+1}$, ζ be a p-th root of unity and $\zeta_j := \zeta^{p_{j+1} \cdots p_{l+1}}$ for $j = 1, \ldots, l$. Then ζ_j is a $p_1 \ldots p_j$-th root of unity for all j, $1 \leq j \leq l$. Define

$$\Psi^{(1)} := \prod_{j=1}^{p_1} (y - \zeta^j \star_p \psi(x)),$$

$$r_1 := \begin{cases} r & \text{if } l = 1, \\ \frac{q_2}{p_1 p_2} & \text{otherwise.} \end{cases}$$

Claim 2.18.1. *There exist monomials* $F_{1j} := b_{1j} x^{\alpha_j^1} y^{\beta_j^1}$ *with* $(\alpha_j^1, \beta_j^1) \in \mathbb{Z}^2$ *and* $0 \leq \beta_j^1 < p_1$ *such that* $q_1 > \delta_0(F_{11}) > \delta_0(F_{12}) > \cdots > \delta_0(F_{1m_1}) > \omega_1 := (p_1 - 1)\frac{q_1}{p_1} + r_1$ *such that*

$$\delta_0(\Psi^{(1)} - (y^{p_1} - a_1^{p_1} x^{q_1} - \sum_{j=1}^{m_1} F_{1j})) \leq \omega_1.$$

Proof. Let

$$\psi_1(x) := a_1 x^{\frac{q_1}{p_1}} + \sum_{j=1}^{k_1} a_{1j} x^{\frac{q_{1j}}{p_1}},$$

$$\tilde{\psi}_1(x, \xi_1) := \psi_1(x) + \xi_1 x^{r_1},$$

and δ_1 be the semidegree defined by

$$\delta_1(f) = \deg_x(f(x, \tilde{\psi}_1(x, \xi_1))).$$

At first note that

$$\tilde{\psi}_1^{p_1} = a_1^{p_1} x^{q_1} + \cdots + p_1 a_1^{p_1 - 1} \xi_1 x^{\omega_1} + \text{l.o.t.}$$

In particular, the x-invariant degree of $\tilde{\psi}_1^{p_1}$ is ω_1. Moreover, writing $c_\omega(\tilde{\psi}_1^{p_1})$ for the coefficient of x^ω in $\tilde{\psi}_1^{p_1}$, we see that for all $\omega \in \mathbb{Q}$, if $c_\omega(\tilde{\psi}_1^{p_1}) \neq 0$, then

1. ω is a $\mathbb{Z}$-linear combination of $\frac{1}{p_1}$ and r_1, and
2. moreover, if $\omega > \omega_1$, then $c_\omega(\tilde{\psi}_1^{p_1}) \in \mathbb{C}$ and $\omega = q/p_1$ for an integer q.

Pick the largest $\omega' > \omega_1$ such that $c_{\omega'}(\tilde{\psi}_1^{p_1}) \neq 0$ and ω' is *not* an integer. Then in particular $\omega' < q_1$. Let $q := \omega' p_1 \in \mathbb{Z}$. Since $\gcd(p_1, q_1) = 1$, there are integers α, β with $0 < \beta < p_1$ such that $q = \alpha p_1 + \beta q_1$. Note that

$$\begin{aligned} \deg_x(x^\alpha \tilde{\psi}_1^\beta) &= \delta_0(x^\alpha y^\beta) = \omega', \text{ and} \\ \deg_{\mathrm{x}}^{\mathrm{inv}}(x^\alpha \tilde{\psi}_1^\beta) &= \alpha + (\beta - 1)\frac{q_1}{p_1} + r_1 = \omega' - \frac{q_1}{p_1} + r_1 < \omega_1 \text{ (since } \omega' < q_1). \end{aligned}$$

Let

$$\tilde{\psi}_{11} := \tilde{\psi}_1^{p_1} - a_1^{p_1} x^{q_1} - \sum_{q_1 > \omega > \omega'} c_\omega(\tilde{\psi}_1^{p_1}) x^\omega - \frac{c'_\omega(\tilde{\psi}_1^{p_1})}{a_1^\beta} x^\alpha \tilde{\psi}_1^\beta.$$

Then $\deg_x(\tilde{\psi}_{11}) < \omega' < q_1 = \deg_x(\tilde{\psi}_1^{p_1})$ and $\deg_{\mathrm{x}}^{\mathrm{inv}}(\tilde{\psi}_{11}) = \omega_1$. Moreover, for all $\omega > \omega_1$, if $c_\omega(\tilde{\psi}_{11}) \neq 0$, then $\omega = q/p_1$ for an integer q. It follows that we can continue the above process as long as the x-degree of the resulting degree-wise Puiseux series is greater than ω_1 and there exist monomials $F_{11}, \ldots, F_{1m_1}$ in x and y with integral exponents such that $\delta_0(F_{1j}) < q_1$, the exponent β_j of y in each F_{1j} satisfies $0 \leq \beta_j < p_1$ and

$$\left.\left(y^{p_1} - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j}\right)\right|_{y=\tilde{\psi}_1} = p_1 a_1^{p_1-1}\xi_1 x^{\omega_1} + \text{l.o.t.} \tag{7}$$

where for all $\omega \in \mathbb{Q}$, if the coefficient of x^ω in the right hand side of (7) is non-zero, then ω is a $\mathbb{Z}$-linear combination of $\frac{1}{p_1}$ and r_1. In particular,

$$\delta_1\left(y^{p_1} - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j}\right) = \deg_x\left(\tilde{\psi}_1^{p_1} - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j}(x, \tilde{\psi}_1(x,\xi_1))\right) = \omega_1 \tag{8}$$

Now

$$\begin{aligned}\Psi^{(1)} &= \prod_{j=1}^{p_1}(y - \zeta^j \star_p \psi(x)) \\ &= \prod_{j=1}^{p_1}(y - \zeta^j \star_p \psi_1(x) + \zeta^j \star_p (\psi_1(x) - \psi(x))).\end{aligned}$$

Since $\deg(\psi_1(x) - \psi(x)) \leq r_1$, it follows that

$$\Psi^{(1)} = \prod_{j=1}^{p_1}(y - \zeta^j \star_p \psi_1(x)) + G \tag{9}$$

for some $G \in \mathbb{C}[y]((x))$ with $\delta_0(G) \leq \omega_1$. Now, remark 2.16 implies that

$$\hat{\Psi}^{(1)} := \prod_{j=1}^{p_1}(y - \zeta^j \star_p \psi_1(x)) = \prod_{j=1}^{p_1}(y - \zeta_1^j \star_{p_1} \psi_1(x))$$

It follows that $\hat{\Psi}^{(1)} \in \mathbb{C}[x, x^{-1}, y]$ (because, e.g. the extension $\mathbb{C}(x^{1/p_1})/\mathbb{C}(x)$ is Galois). Consequently, there is $F_1 \in \mathbb{C}[x, x^{-1}, y]$ such that $\deg_y(F_1) < p_1$, F_1 is a sum of monomials in x and y with δ_0-value greather than ω_1, and

$$\Psi^{(1)} = y^{p_1} - F_1 + \tilde{\Psi}^{(1)} \tag{10}$$

with $\delta_0(\tilde{\Psi}^{(1)}) \leq \omega_1$. Note that $\delta_1(\Psi^{(1)}) = \omega_1$. Therefore identities (8) and (10) imply that

$$\delta_1(F_1 - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j} - \tilde{\Psi}^{(1)}) \leq \omega_1,$$

$$\Rightarrow \delta_1(F_1 - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j}) \leq \omega_1 \text{ (since } \delta_1(\tilde{\Psi}^{(1)}) \leq \delta_0(\tilde{\Psi}^{(1)}) \leq \omega_1).$$

Claim 2.18.2 below then implies that $F_1 = a_1^{p_1}x^{q_1} + \sum_{j=1}^{m_1} F_{1j}$.

Claim 2.18.2. *If F is a δ_0-homogeneous polynomial in $\mathbb{C}[x,y]$ with $\deg_y(F) < p_1$, then $\delta_1(F) = \delta_0(F)$.*

Proof. It is straightforward to see that $\delta_1 \leq \delta_0$. Let I be the δ_0-homogeneous ideal in $\mathbb{C}[x,y]$ generated by δ_0-homogeneous forms f such that $\delta_1(f) < \delta_0(f)$. Then I is the principal ideal generated by $y^{p_1} - a_1^{p_1}x^{q_1}$. Now if $\delta_1(F) < \delta_0(F)$, then the leading form $\mathfrak{L}_{\delta_0}(F)$ is an element of I. Consequently, $y^{p_1} - a_1^{p_1}x^{q_1}$ divides $\mathfrak{L}_{\delta_0}(F)$, which is impossible by the assumption on F. □

□

Claim 2.18.3. *Let $F_{1j} := b_{1j}x^{\alpha_j^1}y^{\beta_j^1}$, $1 \leq j \leq m_1$ be as in claim 2.18.1. Assume there is j, $1 \leq j \leq m_1$, such that α_j^1 is negative. If $\phi_1, \ldots, \phi_m$ are degree-wise Puiseux series in x such that $\phi_j \equiv_{r_1} \psi$ for each j, $1 \leq j \leq m$, then $\prod_{j=1}^m \Phi_j^* \notin \mathbb{C}[x,y]$.*

Proof. Assume to the contrary of the claim that $f := \prod_{j=1}^m \Phi_j^* \in \mathbb{C}[x,y]$. Pick j, $1 \leq j \leq m$. Let s_j denote the polydromy order of ϕ_j and μ_j be a s_j-th root of unity. Note that the first Puiseux pair for ϕ_j is (p_1, q_1). It follows that p_1 divides s_j and

$$\begin{aligned}\Phi_j^{(1)} &:= \prod_{i=1}^{p_1}(y - \mu_j^i \star_{s_j} \phi_j(x)) \\ &= \prod_{i=1}^{p_1}(y - \mu_j^i \star_{s_j} \psi_1(x) + \mu_j^i \star_{s_j} (\psi_1(x) - \phi_j(x))).\end{aligned}$$

Since $\deg(\psi(x) - \phi_j(x)) \leq r_1$, it follows that

$$\Phi_j^{(1)} = \prod_{i=1}^{p_1}(y - \mu_j^i \star_{s_j} \psi_1(x)) + \tilde{G}_j = \prod_{i=1}^{p_1}(y - \zeta_1^i \star_{p_1} \psi_1(x)) + \tilde{G}_j.$$

for some $\tilde{G}_j \in \mathbb{C}[y]((x))$ with $\delta_0(\tilde{G}_j) \leq \omega_1$. Identity (9) then implies that

$$\Phi_j^{(1)} = \psi^{(1)} + \tilde{G}_j - G,$$

where $G \in \mathbb{C}[y]((x))$ and also $\delta_0(G) \leq \omega_1$. Claim 2.18.1 then implies that

$$\Phi_j^{(1)} = y^{p_1} - a_1^{p_1}x^{q_1} - \sum_{i=1}^{m_1} F_{1i} + G_j$$

for some $G_j \in \mathbb{C}[y]((x))$ with $\delta_0(G_j) \leq \omega_1$. Pick the smallest j_0 such that $\alpha_{j_0}^1 < 0$. Let $W_0 := \delta_0(F_{1j_0}) > \omega_1$. Now

$$f = \prod_{i=1}^{M}(y^{p_1} - a_1^{p_1}x^{q_1} - F_{11} - \cdots - F_{1j_0} - \tilde{G}_i)$$

where $\delta_0(\tilde{G}_i) < W_0$ for all i. For each $W \in \mathbb{Q}$, let f_W be the sum of monomials in the right hand side of the above expression with δ_0-value equaling W. Then for all $W > W_1 :=$

$(M-1)q_1+W_0$, f_W is a polynomial in x and y. Moreover, $f_{W_1} = \tilde{F} - M(y^{p_1} - a_1^{p_1}x^{q_1})^{M-1}F_{1j_0}$ for a polynomial $\tilde{F}$ in x and y. It follows that

$$\tilde{f} := f - \sum_{W > W_1} f_W - \tilde{F}$$

is also a polynomial. Since $\mathfrak{L}_{\delta_0}(\tilde{f}) = -M(y^{p_1} - a_1^{p_1}x^{q_1})^{M-1}F_{1j_0}$ is not a polynomial, this is a contradiction and proves the claim. $\square$

Now assume for each j, $1 \leq j \leq m_1$, $\alpha_j^1 \geq 0$. Let $f_1 := y^{p_1} - a_1^{p_1}x^{q_1} - \sum_{j=1}^{m_1} F_{1j}$.

Claim 2.18.4. *$\mathbb{C}[x,y]^{\delta_1}$ is generated as a $\mathbb{C}$-algebra by $(1)_1$, $(x)_1$, $(y)_{q_1/p_1}$ and $(f_1)_{\omega_1}$.*

Proof. Let $A_1 := \mathbb{C}[x,y,y_1]$ where y_1 is an indeterminate and η_1 be the weighted degree on A_1 corresponding to weights 1, q_1/p_1 and ω_1 respectively for x, y and y_1. Let $\pi_1 : A_1 \to A := \mathbb{C}[x,y]$ be the map that sends $y_1 \mapsto f_1$. We have to show that δ_1 is precisely the degree-like function $\bar{\eta}_1$ induced on A by π_1.

Let $J_1 := \mathfrak{L}_{\eta_1}(\ker \pi_1)$, i.e. J_1 is generated by $\{f \in A_1 : f$ is weighted homogeneous (w.r.t. η_1) and $f = \mathfrak{L}_{\eta_1}(g)$ for some $g \in \ker \pi_1\}$. It is straightforward to see that J_1 is the principal ideal generated by $y^{p_1} - a_1^{p_1}x^{q_1}$. Let K_1 be the ideal of A_1 generated by $\{f \in A_1 : f$ is weighted homogeneous (w.r.t. η_1) and $\delta_1(\pi_1(f)) < \eta_1(f)\}$. It suffices to show that $J_1 = K_1$.

Note that K_1 is prime and $K_1 \supseteq J_1$. Also, there is an isomorphism $\chi_1 : A_1/J_1 \cong \mathbb{C}[x, x^{q_1/p_1}, y_1]$ of graded rings, where the gradings on both A_1 and $\mathbb{C}[x,y_1]$ are induced by η_1. Assume to the contrary of the claim that $K_1 \supsetneqq J_1$. Then $\chi_1(K_1)$ is a non-zero homogeneous prime ideal of $\mathbb{C}[x, x^{q_1/p_1}, y_1]$ and therefore $\chi_1(K_1) = \mathfrak{p} \cap \mathbb{C}[x, x^{q_1/p_1}, y_1]$ for a non-zero homogeneous prime ideal $\mathfrak{p}$ of $\mathbb{C}[x^{1/p_1}, y_1]$. Since neither x nor y_1 belongs to $\chi_1(K_1)$, it follows that $\mathfrak{p}$ is generated by an element of the form $y_1^m - cx^{n/q_1}$ for some $c \in \mathbb{C}$ and coprime positive integers m, n such that $\omega_1 = \frac{n}{mq_1}$. But then $y_1^{mq_1} - c^{q_1}x^n \in K_1$, which is impossible by (8). $\square$

Now assume $l \geq 2$ and $\alpha_j^1 \geq 0$ for all j, $1 \leq j \leq m_1$. Let A_1, f_1, δ_1 and η_1 be as in claim 2.18.4. Let us also write y_0 for y. Define

$$r_2 := \begin{cases} r & \text{if } l = 2, \\ \frac{q_3}{p_1p_2p_3} & \text{otherwise.} \end{cases}$$

$$\psi_2(x) := \sum_{\omega > r_2} c_\omega(\psi)x^\omega$$

$$\Psi_2^{(2)} := \prod_{j=1}^{p_1p_2} (y - \zeta_2^j \star_{p_1p_2} \psi_2(x)),$$

Let $\tilde{\psi}_2(x,\xi_2) := \psi_2(x) + \xi_2 x^{r_2}$ and δ_2 be the semidegree defined by

$$\delta_2(f) := \deg_x(f(x, \tilde{\psi}_2(x,\xi_2))).$$

Claim 2.18.5. *There exist monomials $F_{2j} := b_{2j}x^{\alpha_j^2}y^{\beta_{0j}^2}y_1^{\beta_{1j}^2}$, $0 \leq j \leq m_2$ in $\mathbb{C}[x,x^{-1},y,y_1]$ such that*

1. *for each j, $0 \leq j \leq m_2$, $(\alpha^2_j, \beta^2_{0j}, \beta^2_{1j}) \in \mathbb{Z}^3$ and $0 \leq \beta^2_{ij} < p_{i+1}$ for $0 \leq i \leq 1$,*
2. *$\beta^2_{10} = 0$,*
3. *$p_2\omega_1 = \eta_1(F_{20}) > \cdots > \eta_1(F_{2m_2}) > \omega_2 := (p_2 - 1)\omega_1 + (p_1 - 1)\frac{q_1}{p_1} + r_2$,*
4. *$\Psi^{(2)}_2 = f_1^{p_2} - \sum_{j=0}^{m_2} F_{2j}(x, y, f_1) - G_2(x, y, f_1)$ for some $G_2 \in \mathbb{C}[x, x^{-1}, y, y_1]$ such that $\eta_1(G_2) \leq \omega_2$, and*
5. *if $f_2 := f_1^{p_2} - \sum_{j=0}^{m_2} F_{2j}(x, y, f_1) \in \mathbb{C}[x, x^{-1}, y]$, then*

$$\delta_2(f_2) = \deg^{\text{inv}}_{\text{x}}(f_2(x, \tilde{\psi}_2(x, \xi_2)) = \omega_2. \tag{11}$$

Proof. Let ψ_1 and $\tilde{\psi}_1$ be as in claim 2.18.1. Since $l \geq 2$, it follows that $r_1 = \frac{q_2}{p_1 p_2}$ and therefore

$$\begin{aligned} \omega_1 &= (p_1 - 1)\frac{q_1}{p_1} + \frac{q_2}{p_1 p_2}, \text{ and} \\ \psi_2(x) &= \psi_1(x) + x^{r_1}\hat{\psi}_{12}(x), \text{ where} \\ \hat{\psi}_{12}(x) &:= a_2 + \sum_{j=1}^{k_2} a_{2j} x^{-\frac{q_2 - q_{2j}}{p_1 p_2}}. \end{aligned}$$

Recall that $f_1 = y^{p_1} - a_1^{p_1} x^{q_1} - \sum_{j=1}^{m_1} F_{1j} \in \mathbb{C}[x, y, y_1]$. Substituting $\xi_1 = \hat{\psi}_{12}(x) + \xi_2 x^{r_2 - r_1}$ in (7) yields that

$$\begin{aligned} f_1|_{y=\tilde{\psi}_2} &= p_1 a_1^{p_1-1} a_2 x^{\omega_1} + \cdots + p_1 a_1^{p_1-1} \xi_2 x^{\omega_{12}} + \text{l.o.t.}, \text{ where} \\ \omega_{12} &:= (p_1 - 1)\frac{q_1}{p_1} + r_2. \end{aligned}$$

In particular, $\delta_2(f_1) = \omega_1 = \delta_1(f_1)$. Since δ_2 agrees with δ_1 also on x and y, it follows that $\delta_2(H(x, y, f_1)) = \delta_1(H(x, y, y_1)) = \eta_1(H)$ for every monomial H in $A_1 := \mathbb{C}[x, y, y_1]$. The following claim describes one more class of polynomials F in A_1 such that $\delta_2(F(x, y, f_1)) = \eta_1(F)$.

Claim 2.18.6. *If F is a η_1-homogeneous polynomial in A_1 with $\deg_y(F) < p_1$ and $\deg_{y_2}(F) < p_2$, then*

1. *every η_1-homogeneous form of F is a monomial, and*
2. *$\delta_2(F(x, y, y_1)) = \eta_1(F)$.*

Proof. At first note that assertion 1 of the claim implies assertion 2. Indeed, assertion 1 implies that $F = \sum F_i$ such that each F_i is a monomial with $\eta_1(F_1) > \eta_1(F_2) > \cdots$. It follows that $\delta_2(F_1(x, y, y_1)) > \delta_2(F_2(x, y, y_1)) > \cdots$ and consequently $\delta_2(F(x, y, y_1)) = \delta_2(F_1(x, y, y_1)) = \eta_1(F_1) = \eta_1(F)$, which is precisely the content of assertion 2. Therefore it suffices to prove assertion 1.

Let H and H' be two arbitrary distinct monomials that appear in F. It suffices to show that $\eta_1(H) \neq \eta_1(H')$. Let $H = ax^{\alpha}y^{\beta}y_1^{\beta_1}$ and $H' = a'x^{\alpha'}y^{\beta'}y_1^{\beta_1'}$. Assume to the contrary that $\eta_1(H) = \eta_1(H')$. Then

$$\begin{aligned}
&\alpha + \beta\frac{q_1}{p_1} + \beta_1\omega_1 = \alpha' + \beta'\frac{q_1}{p_1} + \beta_1'\omega_1 \\
&\Rightarrow (\beta_1 - \beta_1')((p_1 - 1)\frac{q_1}{p_1} + \frac{q_2}{p_1p_2}) = (\alpha' - \alpha) + (\beta' - \beta)\frac{q_1}{p_1} \\
&\Rightarrow (\beta_1 - \beta_1')q_2 = p_2((\alpha' - \alpha)p_1 + (\beta' - \beta)q_1 + (\beta_1' - \beta_1)(p_1 - 1)q_1).
\end{aligned}$$

Since $\gcd(p_2, q_2) = 1$, it follows that p_2 divides $\beta_1 - \beta_1'$, which implies that $\beta_1 = \beta_1'$ (because $0 \leq \beta_1, \beta_1' < p_2$). Since $\gcd(p_1, q_1) = 1$, this in turn implies that $\beta = \beta'$. Since $\eta_1(H) = \eta_1(H')$, it follows that $\alpha = \alpha'$. This forces H and H' to be the same monomial of F, which is a contradiction. Therefore $\eta_1(H) \neq \eta_1(H')$, as required. $\square$

Let $\psi_{12} := \psi_2 - \psi_1$. Then

$$\begin{aligned}
\Psi_2^{(2)} &= \prod_{i=0}^{p_2-1}\prod_{j=0}^{p_1-1}(y - \zeta_2^{ip_1+j} \star_{p_1p_2} \psi_1(x) - \zeta_2^{ip_1+j} \star_{p_1p_2} \psi_{12}(x)) \\
&= \prod_{i=0}^{p_2-1}\prod_{j=0}^{p_1-1}(y - \zeta_1^{ip_1+j} \star_{p_1} \psi_1(x) - \zeta_2^{ip_1+j} \star_{p_1p_2} \psi_{12}(x)) \quad (\text{since } \zeta_1 = \zeta_2^{p_2}) \\
&= \prod_{i=0}^{p_2-1}\prod_{j=0}^{p_1-1}(y - \zeta_1^{j} \star_{p_1} \psi_1(x) - \zeta_2^{ip_1+j} \star_{p_1p_2} \psi_{12}(x)) \\
&= \prod_{i=0}^{p_2-1}(y - \psi_1(x) - \zeta_2^{ip_1} \star_{p_1p_2} \psi_{12}(x)) \times \prod_{i=0}^{p_2-1}\prod_{j=1}^{p_1-1}(y - \zeta_1^{j} \star_{p_1} \psi_1(x) - \zeta_2^{ip_1+j} \star_{p_1p_2} \psi_{12}(x)).
\end{aligned}$$

Note that the leading term of $\psi_{12}(x)$ is $a_2x^{\frac{q_2}{p_1p_2}}$ and for each $i \geq 0$, the coefficient of $x^{\frac{q_2}{p_1p_2}}$ in $\zeta_2^{ip_1}\star_{p_1p_2}\psi_{12}(x)$ is $a_2\zeta_2^{ip_1q_2}$ which does not equal a_2 unless p_2 divides i (since $\gcd(p_2, q_2) = 1$). It follows that

$$\begin{aligned}
&\deg_x\left(\prod_{i=1}^{p_2-1}(\tilde{\psi}_2(x, \xi_2(x)) - \psi_1(x) - \zeta_2^{ip_1} \star_{p_1p_2} \psi_{12}(x))\right) = (p_2 - 1)\frac{q_2}{p_1p_2}, \text{ and therefore} \\
&\delta_2\left(\Psi_2^{(2)}\right) = \deg_x\left(\Psi_2^{(2)}(x, \tilde{\psi}_2(x, \xi_2(x)))\right) = r_2 + (p_2 - 1)\frac{q_2}{p_1p_2} + p_2(p_1 - 1)\frac{q_1}{p_1} = \omega_2.
\end{aligned}$$

Note also that the leading term of $\Psi_2^{(2)}(x, \tilde{\psi}_2(x, \xi_2(x)))$ is $\tilde{a}_2\xi_2x^{\omega_2}$ for some non-zero $a_2 \in \mathbb{C}$ and therefore

$$\deg_{\mathrm{x}}^{\mathrm{inv}}\left(\Psi_2^{(2)}(x, \tilde{\psi}_2(x, \xi_2(x)))\right) = \delta_2\left(\Psi_2^{(2)}\right) = \omega_2. \tag{12}$$

Claim 2.18.7. *There exists $F \in \mathbb{C}[x, x^{-1}, y, y_1]$ such that*

1. $F(x, y, y_1) = \Psi_2^{(2)}$,
2. $\eta_1(F) = p_2\omega_1$,
3. $\deg_y(F - y_1^{p_2}) < p_1$ *and* $\deg_{y_1}(F - y_1^{p_2}) < p_2$.

Proof. Let $F_0 := \Psi_2^{(2)}$. Then F_0 is an element of $\mathbb{C}[x, x^{-1}, y]$ which is monic in y of degree p_1p_2. We now construct a sequence $\{F_j : j \geq 0\}$ of the elements of $\mathbb{C}[x, x^{-1}, y, y_1]$ as follows:

Assume F_i is given. For every monomial H of F_i, let d_H be the quotient of the division of $\deg_y(H)$ by p_1. Let

$$\tilde{H} := \frac{H}{y^{d_H p_1}}(y_1 + a_1^{p_1} x^{q_1} + \sum_{j=1}^{m_1} F_{1j})^{d_H}$$

and let F_{i+1} be the element of $\mathbb{C}[x, x^{-1}, y, y_1]$ constructed by replacing every monomial H of F_i by $\tilde{H}$.

Note that $\deg_y(F_{i+1}) \leq \deg_y(F_i)$ for every i, and therefore there exists i such that $\deg_y(F_{i'}) = \deg_y(F_i)$ for all $i' \geq i$. It follows that $\deg_y(F_i) < p_1$. Let $F := F_i$. Then it is straightforward to see that assertion 3 of the claim is true. Moreover, the construction of F_j's shows that $F_j(x, y, y_1) = F_{j+1}(x, y, y_1)$ for all $j \geq 0$. In particular, $F(x, y, y_1) = F_0(x, y, y_1) = \Psi_2^{(2)}$, so that assertion 1 is also satisfied. It remains to show that $\eta_1(F) = p_2\omega_1$. Indeed, assume to the contrary that $\eta_1(F) \neq p_2\omega_1$. Since $y_1^{p_2}$ is a monomial of F, it follows that $\eta_1(F) > p_2\omega_1$. In particular, $\deg_{y_1}(\mathfrak{L}_{\eta_1}(F)) < p_2$ and $\deg_y(\mathfrak{L}_{\eta_1}(F)) < p_1$. Then claim 2.18.6 implies that $\delta_2(F(x, y, y_1)) = \eta_1(F) > p_2\omega_1$. On the other hand, $\delta_2(F(x, y, y_1)) = \delta_2(\Psi^{(2)}) = \omega_2 < p_2\omega_1$. This contradiction shows that $\eta_1(F) = p_2\omega_1$ and finishes the proof of the claim. □

Let F be as in claim 2.18.7. Express F as $F = y_1^{p_2} - \sum_{j=0}^{m_2} F_{2j} + G_2$, where $-\sum_{j=0}^{m_2} F_{2j}$ is precisely the sum of monomials H in F with $\eta_1(F) > \omega_2$. An application of claim 2.18.6 to $F - y_1^{p_2}$ shows that $F_{20}, \ldots, F_{2m_2}$ satisfy assertions 1, 3 and 4 of claim 2.18.5. It remains to prove assertions 2 and 5.

Assertion 3 of claim 2.18.5 implies in particular that $\eta_1(F_{20}) = p_2\omega_1$. Note that $p_2\omega_1 \in \frac{1}{p_1}\mathbb{Z}$, and therefore there exists $\alpha, \beta \in \mathbb{Z}$ such that $0 \leq \beta < p_1$ and $\eta_1(x^\alpha y^\beta) = p_2\omega_1$. Assertion 1 of claim 2.18.6 then implies that $F_{20} = ax^\alpha y^\beta$ for some $a \in \mathbb{C}$, which proves assertion 2 of claim 2.18.5.

Let $f_2 := f_1^{p_2} - \sum_{j=0}^{m_2} F_{2j}(x, y, f_1)$ be as in assertion 5 of claim 2.18.5. Recall from (12) that $\delta_2(\Psi^{(2)}) = \omega_2$. Since $\delta_2(H(x, y, f_1)) \leq \eta_1(H)$ for all $H \in A_1$, it follows that

$$\delta_2(f_2) = \delta_2(\Psi_2^{(2)} - G_2(x, y, f_1)) \leq \max\{\delta_2(\Psi_2^{(2)}), \delta_2(G_2)\} = \omega_2.$$

If $\delta_2(f_2) < \omega_2$, then $\delta_2(G_2(x, y, f_1)) = \omega_2 > \delta_2(f_2)$, and therefore claim 2.18.6 implies that there is a monomial H in G_2 such that $\delta_2(H(x, y, f_1)) = \omega_2 > \delta_2(\Psi_2^{(2)} - H(x, y, f_1))$. Now, note that $\deg_{\mathrm{x}}^{\mathrm{inv}}(\tilde{\psi}_2(x, \xi_2(x))) < \deg_x(\tilde{\psi}_2(x, \xi_2(x)))$ and $\deg_{\mathrm{x}}^{\mathrm{inv}}(f_1(x, \tilde{\psi}_2(x, \xi_2(x)))) < \deg_x(f_1(x, \tilde{\psi}_2(x, \xi_2(x))))$. It follows that

$$\deg_{\mathrm{x}}^{\mathrm{inv}}\left(H(x, y, f_1)|_{y=\tilde{\psi}_2(x,\xi_2(x))}\right) < \deg_x\left(H(x, y, f_1)|_{y=\tilde{\psi}_2(x,\xi_2(x))}\right) = \delta_2(H(x, y, f_1)) = \omega_2$$

and consequently $\deg_{\mathrm{x}}^{\mathrm{inv}}(\Psi_2^{(2)}(x, \tilde{\psi}_2(x, \xi_2(x)))) < \omega_2$. But this contradicts (12) and therefore $\delta_2(f_2) = \omega_2$. Moreover, the preceding arguments also show that

$$\deg_{\mathrm{x}}^{\mathrm{inv}}\left(G_2(x, y, f_1)|_{y=\tilde{\psi}_2(x,\xi_2(x))}\right) < \omega_2$$

Therefore, if $\deg_{\mathrm{x}}^{\mathrm{inv}}(f_2(x, \tilde{\psi}_2(x, \xi_2)))$ is also less than ω_2, then it would follow that $\deg_{\mathrm{x}}^{\mathrm{inv}}(\Psi_2^{(2)}(x, \tilde{\psi}_2(x, \xi_2(x))))$ is less than ω_2, which leads to the same contradiction. Consequently $\deg_{\mathrm{x}}^{\mathrm{inv}}(f_2(x, \tilde{\psi}_2(x, \xi_2))) = \omega_2$. This proves (11) and finishes the proof of claim 2.18.5. □

Claim 2.18.8. *Let* $F_{2j} := b_{2j} x^{\alpha^2_j} y^{\beta^2_{0j}} y_1^{\beta^2_{1j}} \in A_1$, $0 \leq j \leq m_2$ *be as in claim 2.18.5. Assume there is* j, $0 \leq j \leq m_2$, *such that* α^2_j *is negative. If* $\phi_1, \ldots, \phi_m$ *are degree-wise Puiseux series in* x *such that* $\phi_j \equiv_{r_2} \psi$ *for each* j, $1 \leq j \leq m$, *then* $\prod_{j=1}^m \Phi_j^* \notin \mathbb{C}[x, y]$.

Proof. Assume to the contrary of the claim that $f := \prod_{j=1}^m \Phi_j^* \in \mathbb{C}[x, y]$. Pick j, $1 \leq j \leq m_2$. Let s_j denote the polydromy order of ϕ_j and μ_j be a s_j-th root of unity. Note that the first two Puiseux pairs for ϕ_j are (p_1, q_1) and (p_2, q_2). It follows that $p_1 p_2$ divides s_j and

$$\begin{aligned}
\Phi_j^{(1)} &:= \prod_{i=1}^{p_1} (y - \mu_j^i \star_{s_j} \phi_j(x)) \\
&= \prod_{i=1}^{p_1} (y - \mu_j^i \star_{s_j} \psi_2(x) + \mu_j^i \star_{s_j} (\psi_2(x) - \phi_j(x))).
\end{aligned}$$

Since $\deg(\psi(x) - \phi_j(x)) \leq r_2$, it follows that

$$\begin{aligned}
\Phi_j^{(1)} &= \prod_{i=1}^{p_1} (y - \mu_j^i \star_{s_j} \psi_2(x)) + \tilde{G}_j \ \text{(where } \tilde{G}_j \in \mathbb{C}[y]((x)),\ \delta_0(\tilde{G}_j) \leq \omega_{12}) \\
&= \prod_{i=1}^{p_1} (y - \zeta_2^i \star_{p_1 p_2} \psi_2(x)) + \tilde{G}_j \\
&= f_1 + g_{12} + \hat{G}_j \ \text{(where } g_{12} := \sum_{\omega_1 \geq \delta_0(x^\alpha y^\beta) > \omega_{12}} c_{\alpha,\beta}(\Psi_2^{(1)}) x^\omega \text{ and } \delta_0(\hat{G}_j) \leq \omega_{12})
\end{aligned}$$

Let $G_j^{(1)} := y_1 + g_{12}(x, y) + \hat{G}_j(x, y)$. Then

$$\begin{aligned}
\Phi_j^{(2)} &= \prod_{i=1}^{p_2} \mu_j^{ip_1} \star_{s_j} \Phi_j^{(1)} = G_j^{(2)}(x, y, f_1), \text{ where} \\
G_j^{(2)} &:= \prod_{i=1}^{p_2} \mu_j^{ip_1} \star_{s_j} G_j^{(1)} = \prod_{i=1}^{p_2} (y_1 + \mu_j^{ip_1} \star_{s_j} g_{12} + \mu_j^{ip_1} \star_{s_j} \hat{G}_j) \\
&= \prod_{i=1}^{p_2} (y_1 + \zeta_2^{ip_1} \star_{p_1 p_2} g_{12} + \mu_j^{ip_1} \star_{s_j} \hat{G}_j) \\
&= y_1^{p_2} - F_2(x, y, y_1) + H_j(x, y, y_1),
\end{aligned}$$

where $F_2 := \sum_{j=0}^{m_2} F_{2j}$ and $H_j \in \mathbb{C}[x^{\frac{1}{s_j}}, x^{-\frac{1}{s_j}}, y, y_1]$ such that $\eta_1(H_j) \leq \omega_2$. Let $t_j := \frac{s_j}{p_1 p_2}$. Then t_j is a positive integer and

$$\Phi_j^* = \prod_{i=0}^{t_j - 1} \prod_{k=0}^{p_1 p_2 - 1} \mu_j^{ip_1 p_2 + j} \star_{s_j} (y - \phi_j(x)) = \prod_{i=0}^{t_j - 1} \mu_j^{ip_1 p_2} \star_{s_j} \Phi_j^{(2)} = G_j^*(x, y, f_1) \text{ where}$$

$$G_j^* := \prod_{i=0}^{t_j - 1} \mu_j^{ip_1 p_2} \star_{s_j} G_j^{(2)} = \prod_{i=0}^{t_j - 1} \prod_{k=0}^{p_2 - 1} \mu_j^{ip_1 p_2 + kp_1} \star_{s_j} G_j^{(1)} = \prod_{i=0}^{p_2 t_j - 1} \mu_j^{ip_1} \star_{s_j} G_j^{(1)}$$

Since Φ_j^* is invariant under the action of μ_j, it follows that $G_j^{**}(x, y, y_1) = (\Phi_j^*)^{p_1}$, where

$$G_j^{**} := \prod_{k=0}^{p_1-1} \mu_j^k \star_{s_j} G_j^* = \prod_{k=0}^{p_1-1} \prod_{i=0}^{p_2 t_j - 1} \mu_j^{ip_1+k} \star_{s_j} G_j^{(1)} = \prod_{i=0}^{s_j-1} \mu_j^i \star_{s_j} G_j^{(1)}.$$

The above expression implies that G_j^{**} is invariant under the action of μ_j. Since the polydromy order of G_j^{**} is s_j, it follows that all exponents of x in the expansion of G_j^{**} are integral. Let

$$\begin{aligned} G^{**} &:= \prod_{j=1}^{m} G_j^{**} = \prod_{j=1}^{m} \prod_{k=0}^{p_1-1} \mu_j^k \star_{s_j} G_j^* = \prod_{j=1}^{m} \prod_{k=0}^{p_1-1} \prod_{i=0}^{t_j-1} \mu_j^{ip_1p_2+k} \star_{s_j} G_j^{(2)} \\ &= \prod_{j=1}^{m} \prod_{k=0}^{p_1-1} \prod_{i=0}^{t_j-1} (y_1^{p_2} - F_2(x, y, y_1) + \mu_j^{ip_1p_2+k} \star_{s_j} H_j(x, y, y_1)). \end{aligned}$$

Pick the smallest j_0 such that $\alpha^2_{j_0} < 0$. Let $W_0 := \eta_1(F_{2j_0})$. Note that $W_0 > \omega_2$ and therefore

$$G^{**} = \prod_{i=1}^{M} (y_1^{p_2} - F_{20}(x, y, y_1) - \cdots - F_{2j_0}(x, y, y_1) + \tilde{H}_i(x, y, y_1))$$

where $M := mp_1t_j$ and $\eta_1(\tilde{H}_i) < W_0$ for all i. Let

$$G^* := \begin{cases} G^{**} - y_1^{Mp_2} & \text{if } j_0 = 1, \\ G^{**} - (y_1^{p_2} - F_{20}(x, y, y_1) - \cdots - F_{2,j_0-1}(x, y, y_1))^M & \text{otherwise.} \end{cases}$$

Then $\pi_1(G^*) \in \mathbb{C}[x, y]$ (recall that $\pi_1 : \mathbb{C}[x, y, y_1] \to \mathbb{C}[x, y]$ is identity on $\mathbb{C}[x, y]$ and maps $y_1 \mapsto f_1$) and

$$\mathfrak{L}_{\eta_1}(G^*) = \begin{cases} -F_{20} & \text{if } j_0 = 0 \text{ and } M = 1, \\ -My_1^{(M-1)p_2} F_{20} + \sum_{i=2}^{M} \binom{M}{i} y_1^{(M-i)p_2} (-F_{20})^i & \text{if } j_0 = 0, \text{and } M > 1, \\ -M(y_1^{p_2} - F_{20})^{M-1} F_{2,j_0} & \text{otherwise.} \end{cases}$$

Let $f^* := \pi_1(G^*) \in \mathbb{C}[x, y]$. Since δ_1 is induced by η_1, there is a polynomial $F^* \in A_1 := \mathbb{C}[x, y, y_1]$ such that $\pi_1(F^*) = f^*$ and $\eta_1(F^*) = \delta_1(f^*) \leq \eta_1(G^*)$ (where the last inequality holds because $f^* = \pi_1(G^*)$).

Claim 2.18.9. *$\eta_1(F^*) = \eta_1(G^*)$.*

Proof. Let $v_1 := (\eta(x), \eta(y), \eta(f_1))$ and Define an ordering $\prec_1$ on $\mathbb{Z}^3_{\geq 0}$ by setting $(\alpha, \beta, \beta_1) \prec_1 (\alpha', \beta', \beta_1')$ iff

1. $(\alpha - \alpha', \beta - \beta', \beta_1 - \beta_1') \cdot v_1 < 0$, or
2. $(\alpha - \alpha', \beta - \beta', \beta_1 - \beta_1') \cdot v_1 = 0$ and the right-most non-zero entry of $(\alpha - \alpha', \beta - \beta', \beta_1 - \beta_1')$ is negative.

Then $\prec_1$ induces a monomial ordering on A_1. Let K_1 be the ideal of A_1 generated by $\{f \in A_1 : f \text{ is weighted homogeneous (w.r.t. } \eta_1) \text{ and } \delta_1(\pi_1(f)) < \eta_1(f)\}$. Recall that we proved in claim 2.18.4 that K_1 is the principal ideal generated by $h_1 := y^{p_1} - a_1^{p_1} x^{q_1}$. It follows that the Gröbner basis of K_1 with respect to $\prec_1$ is $\mathcal{B}_1 = \{h_1\}$ and the leading term of

h_1 with respect to $\prec_1$ is $\mathrm{LT}_{\prec_1}(h_1) = y^{p_1}$.

Assume to the contrary of the claim that $\eta_1(F^*) < \eta_1(G^*)$. Then

$$\delta_1(\pi_1(\mathfrak{L}_{\eta_1}(G^*))) = \delta_1(\pi_1(F^*) - \pi_1(G^* - \mathfrak{L}_{\eta_1}(G^*))) < \eta_1(G^*).$$

Hence, if n is the smallest positive integer such that $x^n\mathfrak{L}_{\eta_1}(G^*) \in \mathbb{C}[x, y]$, then $x^n\mathfrak{L}_{\eta_1}(G^*) \in K_1$ and therefore $y^{p_1} = \mathrm{LT}_{\prec_1}(h_1)$ divides $\mathrm{LT}_{\prec_1}(x^n\mathfrak{L}_{\eta_1}(G^*)) = \mathrm{LT}_{\prec_1}(\mathfrak{L}_{\eta_1}(x^nG^*))$. Recall (from assertion 2 of claim 2.18.5) that $\beta^2_{10} = 0$, and therefore

$$\mathrm{LT}_{\prec_1}(\mathfrak{L}_{\eta_1}(x^nG^*)) = \begin{cases} -b_{10}y^{\beta^2_{00}} & \text{if } j_0 = 0 \text{ and } M = 1, \\ -Mb_{10}x^{(M-1)|\alpha^2_0|}y^{\beta^2_{00}}y_1^{(M-1)p_2} & \text{if } j_0 = 0, \text{and } M > 1, \\ -Mb_{1j_0}y^{\beta^2_{0j_0}}y_1^{(M-1)p_2+\beta^2_{1j_0}} & \text{otherwise.} \end{cases}$$

Since $\beta^2_{0j} < p_1$ for all $j \geq 0$, it follows that y^{p_1} does *not* divide $\mathrm{LT}_{\prec_1}(\mathfrak{L}_{\eta_1}(x^nG^*))$ in any case, which is a contradiction. This completes the proof of the claim. ☐

Claim 2.18.9 implies that $\eta_1(F*) = \eta_1(G*)$ and therefore $H^* := \mathfrak{L}_{\eta_1}(x^nG^*) - \mathfrak{L}_{\eta_1}(x^nF^*) \in K_1$, where n is as in claim 2.18.9. Let

$$H_1^* := \mathrm{LT}_{\prec_1}(\mathfrak{L}_{\eta_1}(x^nG^*)) \text{ and } H_2^* := \mathrm{LT}_{\prec_1}(\mathfrak{L}_{\eta_1}(x^nF^*)).$$

Now, note that by our choice of F^*, $\mathfrak{L}_{\eta_1}(x^nF^*) \notin J_1 = K_1$. Therefore, 'dividing out' by $\mathcal{B}_1$ (which does not change $\eta_1(F^*)$), we may assume that y^{p_1} does *not* divide H_2^*. Since $F^* \in \mathbb{C}[x, y]$, it follows that

$$\deg_x(H_2^*) \geq n > \deg_x(H_1^*).$$

In particular, $H_1^* \neq H_2^*$ and

$$\mathrm{LT}_{\prec_1}(H^*) = \max\{H_1^*, H_2^*\}.$$

Since we have shown in the proof of claim 2.18.9 that $\deg_y(H_1^*) < p_1$, it follows that y^{p_1} does not divide $\mathrm{LT}_{\prec_1}(H^*)$. A Gröbner basis argument as in the proof of claim 2.18.9 then implies that $H^* \notin K_1$, which is a contradiction. This completes the proof of the claim. ☐

Now assume that $\alpha^2_j \geq 0$ for each j, $0 \leq j \leq m_2$. Let $F_2 := y_1^{p_2} - \sum_{j=0}^{m_2} F_{2j} \in A_1$. Let y_2 be a new indeterminate, $A_2 := A_1[y_2] = \mathbb{C}[x, y, y_1, y_2]$ and π_2 be the $\mathbb{C}$-algebra homomorphism $A_2 \to \mathbb{C}[x, y]$ which extends π_1 and maps y_2 to $f_2 := F_2(x, y, f_1)$. Finally, let η_2 be the weighted degree on A_2 which extends η_1 and satisfies: $\eta_2(y_2) = \omega_2 = \delta_2(f_2)$.

Claim 2.18.10. *δ_2 is precisely the degree-like function on $\mathbb{C}[x, y]$ induced by η_2 (via π_2).*

Proof. Let J_2 be the ideal of A_2 generated by $h_1 := y^{p_1} - a_1^{p_1}x^{q_1}$ and $h_2 := y_1^{p_2} - F_{20}$. At first we show that J_2 is a prime ideal and determine the quotient A_2/J_2.

Let $\tilde{\chi}_1 : A_1 \to \mathbb{C}[x_1, y_1] =: B_1$ be the $\mathbb{C}$-algebra homomorphism that maps $x \mapsto x_1^{p_1}$, $y \mapsto a_1x_1^{q_1}$ and $y_1 \mapsto y_1$. Then $\ker\tilde{\chi}_1 = \langle h_1\rangle$ and $\tilde{\chi}_1$ induces an isomorphism of $A_1/\langle h_1\rangle$ onto the subring $\tilde{B}_1 := \mathbb{C}[x_1^{p_1}, x_1^{q_1}, y_1]$ of B_1. Recall from claim 2.18.5 that $F_{20} = b_{20}x^{\alpha^2_0}y^{\beta^2_{00}}$ with $c \in \mathbb{C}$ and $\eta_2(x^{\alpha^2_0}y^{\beta^2_{00}}) = p_2\omega_1$. Therefore $\tilde{\chi}_1(h_2) = y_1^{p_2} - b_{20}a_1^{\beta^2_{00}}x_1^{p_1p_2\omega_1}$. Note that $p_1p_2\omega_1 = (p_1 - 1)p_2q_1 + q_2$ which is relatively prime to p_2. Therefore $\tilde{\chi}_1(h_2)$ is an irreducible element of B_1 and the ideal L_1 in B_1 generated by $\tilde{\chi}_1(h_2)$ is prime. We claim that $\tilde{\chi}_1(J_2 \cap A_1) = L_1 \cap \tilde{B}_1$. Indeed, $\tilde{\chi}_1(J_2 \cap A_1)$ is precisely the ideal generated by $\tilde{\chi}_1(h_2)$ in $\tilde{B}_1$ and therefore $\tilde{\chi}_1(J_2 \cap A_1) \subseteq L_1 \cap \tilde{B}_1$. To see the inclusion in the other direction, consider B_1 as a graded $\mathbb{C}$-algebra with the grading given by the weighted degree induced by η_2 (i.e. the weighted degree which corresponds to weights $1/p_1$ for x_1 and ω_1 for y_1). Then $\tilde{B}_1$ is a graded sub-algebra of B_1 and $\tilde{\chi}_1(h_2)$ is a homogeneous element of $\tilde{B}_1$. Since L_1 is also a homogeneous ideal of B_1, it suffices to show that every homogeneous element of $L_1 \cap B_1$ belongs to $\tilde{\chi}_1(J_2 \cap A_1)$. Let h be a non-zero homogeneous element of $L_1 \cap B_1$. Then

$$h = a x_1^{\alpha} y_1^{\beta} (y_1^{p_2} - b_{20} a_1^{\beta_{00}^2} x_1^{p_1 p_2 \omega_1}) \prod_{j=1}^{k} (y_1^{p_2} - c_j x_1^{p_1 p_2 \omega_1})$$

for non-negative integers α, β and non-zero complex numbers $a, c_1, \ldots, c_m$. Note that the coefficient of $x_1^{\alpha} y_1^{(k+1)p_2+\beta}$ in h is non-zero. Since $h \in \tilde{B}_1$, it follows that $x_1^{\alpha} \in \tilde{B}_1$. Moreover, $x_1^{p_1 p_2 \omega_1} = a_1^{-\beta_{00}^2} \tilde{\chi}_1(x^{\alpha_0^2} y^{\beta_{00}^2}) \in \tilde{B}_1$. It follows that h is in the ideal generated by $\tilde{\chi}_1(h_1) = y_1^{p_2} - b_{20} a_1^{\beta_{00}^2} x_1^{p_1 p_2 \omega_1}$ in $\tilde{B}_1$. This completes the proof for the claim that $\tilde{\chi}_1(J_2 \cap A_1) = L_1 \cap \tilde{B}_1$. In particular, it follows that $\chi_1(J_2 \cap A_1)$ is a prime ideal of $\tilde{B}_1$ and therefore $J_2 \cap A_1$ is a prime ideal of A_1. Moreover, there is a chain of graded ring homomorphisms as follows

$$A_1/(J_2 \cap A_1) \cong \tilde{B}_1/(L_1 \cap \tilde{B}_1) \hookrightarrow B_1/L_1 \cong \mathbb{C}[x_1, x_1^{p_1 \omega_1}] \cong \mathbb{C}[x^{\frac{1}{p_1}}, x^{\omega_1}] \hookrightarrow \mathbb{C}[x^{\frac{1}{p_1 p_2}}].$$

Since $A_2/J_2 \cong A_1/(J_2 \cap A_1)[y_2]$, it follows that J_2 is a prime ideal of A_2 and there is a graded ring homomorphism $\chi_2 : A_2 \to \mathbb{C}[x^{\frac{1}{p_1 p_2}}, y_2] =: B_2$ such that $\ker \chi_2 = J_2$. Moreover, since B_1 is integral over $\tilde{B}_1$ and $\mathbb{C}[x^{\frac{1}{p_1 p_2}}]$ is integral over $\mathbb{C}[x^{\frac{1}{p_1}}, x^{\omega_1}]$, it follows that B_2 is integral over $\chi_2(A_2)$. Let

$$\begin{aligned} K_2 := \{f \in A_2 : f = \sum_{j=1}^{k} f_j,\ f_j \text{ is weighted homogeneous w.r.t. } \eta_2 \\ \text{and } \delta_2(\pi_2(f_j)) < \eta_2(f_j) \text{ for each } j,\ 1 \leq j \leq k, \}. \end{aligned}$$

Since $\delta_2(\pi_2(f)) \leq \eta_2(f)$ for all $f \in A_2$, it follows that K_2 is a prime ideal of A_2. We claim that $K_2 = J_2$. Indeed, both h_1 and h_2 are elements of K_2, and therefore $K_2 \supseteq J_2$. Assume contrary to the claim that $K_2 \supsetneq J_2$. Since K_2 is a homogeneous ideal (with respect to the grading induced by η_2), it follows that $\chi_2(K_2)$ is a non-zero homogeneous prime ideal in $\chi_2(A_2)$. Since B_2 is integral over $\chi_2(A_2)$, there exists a non-zero homogeneous prime ideal L_2 of B_2 such that $\chi_2(A_2) = L_2 \cap \chi_2(A_2)$. Since L_2 is homogeneous, either $x^{\frac{1}{p_1 p_2}} \in L_2$, or $y_2 \in L_2$, or $y_2 - c x^{\omega_2} \in L_2$ for some non-zero $c \in \mathbb{C}$. It follows that either $x \in K_2$, or $y_2 \in K_2$, or $y_2^p - a x^p \in K_2$ for some non-zero $a \in \mathbb{C}$ and positive integers p, q such that $q/p = \omega_2$. It follows straight from the definition of K_2 and η_2 that the first two cases are impossible. The last case is impossible due to (11). Therefore $K_2 = J_2$.

Let $\mathfrak{L}_{\eta_2}(\ker \pi_2)$ be the ideal in A_2 generated by all weighted homogeneous (w.r.t. η_2) forms $f \in A_2$ such that $f = \mathfrak{L}_{\eta_2}(g)$ for some $g \in \ker \pi_2$. It follows from the definition that $J_2 \subseteq \mathfrak{L}_{\eta_2}(\ker \pi_2) \subseteq K_2$. Since $J_2 = K_2$, it follows that $\mathfrak{L}_{\eta_2}(\ker \pi_2) = K_2$. This implies that for all $f \in A$,

$$\delta_2(f) = \min\{\eta_2(g) : g \in A_2,\ \pi_2(g) = f\}.$$

In other words, δ_2 is precisely the degree-like function on $\mathbb{C}[x, y]$ induced by η_2, as required. $\square$

We now begin the proof for the general case. At first we describe the set-up for induction. Let

$$r_k := \begin{cases} \frac{q_k}{p_1 p_2 \cdots p_k} & \text{if } 1 \leq k \leq l, \\ r & \text{if } k = l+1, \end{cases}$$

For each k, $0 \leq k \leq l$, let ξ_k be an indeterminate, ζ_k be a $(p_1 \ldots p_k)$-th root of unity (for $k = 0$, we define $\zeta_0 := 1$ and $p_0 := 1$), and

$$\psi_k(x) := \sum_{\omega > r_{k+1}} c_\omega(\psi) x^\omega, \qquad \tilde{\psi}_k(x, \xi_k) := \psi_k(x) + \xi_k x^{r_{k+1}},$$

$$\Psi_k^*(x, y) := \prod_{j=1}^{p_0 \cdots p_k} (y - \zeta_k^j \star_{p_0 \cdots p_k} \psi_k(x)),$$

$$\delta_k := \text{the semidegree on } \mathbb{C}[x, y] \text{ defined by}$$

$$\delta_k(f) := \deg_x(f(x, \tilde{\psi}_k(x, \xi_k))) \text{ for all } f \in \mathbb{C}[x, y],$$

$$\omega_k := \sum_{j=1}^{k} \hat{\sigma}_{k,j} + r_{k+1}, \text{ where for all } m, n \in \mathbb{Z} \text{ such that } 1 \leq n \leq m \leq l+1,$$

$$\hat{\sigma}_{m,n} := \sum_{1 \leq i_1 < \ldots < i_n \leq m} (p_{i_n} - 1)(p_{i_{n-1}} - 1) \cdots (p_{i_1} - 1) r_{i_1},$$

Lemma 5.5 shows that for all j, $0 \leq j \leq l$,

$$\delta_j \left(\Psi_j^*\right) = \deg_{\mathrm{x}}^{\mathrm{inv}} \left(\Psi_j^*(x, \tilde{\psi}_j(x, \xi_j))\right) = \omega_j, \text{ and moreover,} \tag{13}$$

$$\delta_k \left(\Psi_j^*\right) = \omega_j \text{ for all } k \text{ such that } j \leq k \leq l, \tag{14}$$

Let $y_0, \ldots, y_l$ be indeterminates. For each k, $0 \leq k \leq l$, let $A_k := \mathbb{C}[x, y_0, \ldots, y_k]$. Let η be the weighted degree on A_l corresponding to weight 1 for x and ω_j for y_j for $0 \leq k \leq l$. For each k, $0 \leq k \leq l$, let η_k be the restriction of η to A_k. In particular, for all j, k such that $0 \leq j \leq k \leq l$, η_j and η_k agrees on A_j.

Now fix j, $1 \leq j < l$. Assume that there is a sequence of polynomials $F_1, \ldots, F_j$ such that for each k, $1 \leq k \leq j$,

$0_{(k)}$. $F_k \in A_{k-1}$,

$1_{(k)}$. $F_k = y_{k-1}^{p_k} + \sum_{i=0}^{m_k} F_{k,i}$ for some $m_k \geq 0$, where each $F_{k,i}$ is a *monomial* in $x, y_0, \ldots, y_{k-1}$,

$2_{(k)}$. $\deg_{y_{k-1}}(F_{k,0}) = 0$,

$3_{(k)}$. $\deg_{y_s}(F_{k,i}) < p_{s+1}$ for all i, s, $0 \leq i \leq m_k$, $0 \leq s < k$, and

$4_{(k)}$. $p_k \omega_{k-1} = \eta(F_{k,0}) > \cdots > \eta(F_{k,m_k}) > \omega_k$.

Define

$$f_k := \begin{cases} y & \text{if } k = 0, \\ F_k(x, f_0, \ldots, f_{k-1}) & \text{if } 1 \leq k \leq j. \end{cases}$$

Assume in addition that for all k, $1 \leq k \leq j$,

$5_{(k)}$. f_k is monic in y with $\deg_y(f_k) = p_1 p_2 \cdots p_k$,

$6_{(k)}$. $\Psi_k^* = f_k + G_k(x, f_0, \dots, f_{k-1})$ for some $G_k \in A_{k-1}[x^{-1}]$ such that $\eta(G_k) \leq \omega_k$,

$7_{(k)}$. $f_k(x, \tilde{\psi}_k(x, \xi_k)) = c_k \xi_k x^{\omega_k} +$ l.o.t. for some non-zero $c_k \in \mathbb{C}$, and ω is a $\mathbb{Z}$-linear combination of $1, r_1, \dots, r_{k+1}$ for all ω such that the coefficient of x^ω in $f_k(x, \tilde{\psi}_k(x, \xi_k))$ is non-zero. In particular, $\delta(f_k) = \delta_k(f_k) = \deg_{\mathrm{x}}^{\mathrm{inv}}(f_k(x, \tilde{\psi}_k(x, \xi_k))) = \omega_k$.

$8_{(k)}$. δ_k is precisely the degree-like function on $\mathbb{C}[x, y]$ induced by η_k via π_k, where $\pi_k : A_k \to \mathbb{C}[x, y]$ is the $\mathbb{C}$-algebra homomorphism that sends $x \mapsto x$ and $y_i \mapsto f_i$ for $0 \leq i \leq k$.

For each k, $1 \leq k \leq l$, let $v_k := (1, \omega_0, \dots, \omega_k)$ and $\prec_k$ be the ordering on $\mathbb{Z}_{\geq 0}^{k+2}$ by setting $(\alpha, \beta_0, \dots, \beta_k) \prec (\alpha', \beta'_0, \dots, \beta'_k)$ iff

1. $(\alpha - \alpha', \beta_0 - \beta'_0, \dots, \beta_k - \beta'_k) \cdot v_k < 0$, or
2. $(\alpha - \alpha', \beta_0 - \beta'_0, \dots, \beta_k - \beta'_k) \cdot v_k = 0$ and the right-most non-zero entry of $(\alpha - \alpha', \beta_0 - \beta'_0, \dots, \beta_k - \beta'_k)$ is negative.

It follows from its definition that $\prec_k$ is a total ordering on $\mathbb{Z}_{\geq 0}^{k+2}$ and $\prec_k$ is compatible with addition.

Since δ is positive on non-constant polynomials in x and y, assertions $5_{(k)}$ and $7_{(k)}$ imply that $\omega_k > 0$ for all k, $1 \leq k \leq j$. Therefore $\prec_k$ induces a monomial ordering on A_k for all k, $1 \leq k \leq j$. Assume that for all k, $1 \leq k \leq j$,

$9_{(k)}$. Let J_k be the ideal in A_k generated by all weighted homogeneous (w.r.t. η_k) polynomials f such that $\delta_k(f(x, f_0, \dots, f_k)) < \eta_k(f)$. Then $\mathcal{B}_k := (y_{k-1}^{p_k} + F_{k,0}, y_{k-2}^{p_{k-1}} + F_{k-1,0}, \dots, y_0^{p_1} + F_{1,0})$ is a Gröbner basis of J_k with respect to $\prec_k$.

$10_{(k)}$. There exists an isomorphism $\chi_k : A_k/J_k \cong R_k \subseteq \mathbb{C}[x_k, y_k] =: S_k$ such that χ_k sends $x \mapsto x_k^{p_1 \cdots p_k}$, $y_k \mapsto y_k$, and $y_i \mapsto b_i x_k^{p_1 \cdots p_k \omega_i}$ for some $b_i \neq 0 \in \mathbb{C}$ for each i, $0 \leq i \leq k-1$. Consider A_k and S_k as graded rings, the grading on A_k being induced by η_k and the grading on S_k being induced by the weighted degree with weights $\frac{1}{p_1 \cdots p_k}$ for x_k and ω_k for y_k. Then R_k is a graded subring of S_k and χ_k is an isomorphism of graded rings.

Remark 2.19. Assertions $9_{(k)}$ and $10_{(k)}$ are in fact consequences of assertion $8_{(k)}$. We nonetheless list them separately because this simplifies the proof of $8_{(j+1)}$.

We now show that the sequence $\{F_1, \dots, F_j\}$ can be extended by an element F_{j+1} provided we allow the possiblity of negative exponents of x, i.e. F_{j+1} will be an element of $A_j[x^{-1}]$.

Claim 2.19.1. *There is an element $F_{j+1} \in A_j[x^{-1}]$ such that*

$1_{(j+1)}$. $F_{j+1} = y_j^{p_{j+1}} + \sum_{i=0}^{m_{j+1}} F_{j+1,i}$ *for some $m_{j+1} \geq 0$, where each $F_{j+1,i}$ is a* monomial *in $x, y_0, \dots, y_j$.*

$2_{(j+1)}$. $\deg_{y_j}(F_{j+1,0}) = 0$.

$3_{(j+1)}$. $\deg_{y_k}(F_{j+1,i}) < p_{k+1}$ *for all i, k, $0 \leq i \leq m_{j+1}$, $0 \leq k \leq j$.*

$4_{(j+1)}$. $p_{j+1}\omega_j = \eta(F_{j+1,0}) > \cdots > \eta(F_{j+1,m_{j+1}}) > \omega_{j+1}$.

$5_{(j+1)}$. *Define* $f_{j+1} := F_{j+1}(x, f_0, \ldots, f_j)$. *Then* f_{j+1} *is monic in* y *with* $\deg_y(f_{j+1}) = p_1 p_2 \cdots p_{j+1}$.

$6_{(j+1)}$. $\Psi^*_{j+1} = f_{j+1} + G_{j+1}(x, f_0, \ldots, f_j)$ *for some* $G_{j+1} \in A_{j+1}[x^{-1}]$ *such that* $\eta(G_{j+1}) \leq \omega_{j+1}$.

$7_{(j+1)}$. $f_{j+1}(x, \tilde{\psi}_{j+1}(x, \xi_{j+1})) = c_{j+1}\xi_{j+1}x^{\omega_{j+1}} +$ *l.o.t. for some non-zero* $c_{j+1} \in \mathbb{C}$, *and* ω *is a* $\mathbb{Z}$*-linear combination of* $1, r_1, \ldots, r_{j+2}$ *for all* ω *such that the coefficient of* x^ω *in* $f_{j+1}(x, \tilde{\psi}_{j+1}(x, \xi_{j+1}))$ *is non-zero. In particular,* $\delta(f_{j+1}) = \delta_{j+1}(f_{j+1}) = \deg_{\mathrm{x}}^{\mathrm{inv}}(f_{j+1}(x, \tilde{\psi}_{j+1}(x, \xi_{j+1}))) = \omega_{j+1}$.

Proof. Fix k, $1 \leq k \leq j$. The leading term of $\psi_{j+1}(x) - \psi_k(x)$ is $a_{k+1}x^{r_{k+1}}$ and therefore

$$\tilde{\psi}_{j+1}(x, \xi_{j+1}) = \tilde{\psi}_k(x, \xi_k)\Big|_{\xi_k = a_{k+1} + \cdots + \xi_{j+1}x^{r_{j+2} - r_{k+1}}}.$$

Consequently, asertion $7_{(k)}$ implies that

$$f_k(x, \tilde{\psi}_{j+1}(x, \xi_{j+1})) = c_k a_{k+1} x^{\omega_k} + \cdots + c_k \xi_{j+1} x^{\omega_k + r_{j+2} - r_{k+1}} + \text{l.o.t.}, \text{ so that} \tag{15}$$

$$\delta_{j+1}(f_k) = \omega_k = \delta_k(f_k), \text{ and } \deg_{\mathrm{x}}^{\mathrm{inv}}\left(f_k|_{y = \tilde{\psi}_{j+1}(x, \xi_{j+1})}\right) = \omega_k - (r_{k+1} - r_{j+2}) =: \omega_{k,j+2}. \tag{16}$$

Moreover, inductive assumption $7_{(k)}$ implies that

$$c_\omega\left(f_k|_{y = \tilde{\psi}_{j+1}}\right) \neq 0 \quad \Rightarrow \quad \omega \in \langle 1, r_1, \ldots, r_{j+2} \rangle, \tag{17}$$

$$c_\omega\left(f_k|_{y = \tilde{\psi}_{j+1}}\right) \neq 0 \text{ and } \omega > \omega_{k,j+2} \quad \Rightarrow \quad \omega \in \langle 1, r_1, \ldots, r_{j+1} \rangle = \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}. \tag{18}$$

Note that $\omega_{j+1} = p_{j+1}\omega_j + r_{j+2} - r_{j+1}$ (according to assertion 1 of lemma 5.6). Therefore, setting $k = j$ in (15) and taking p_{j+1}-th power yields that

$$\left(f_j|_{y = \psi_{j+1}}\right)^{p_{j+1}} = \tilde{c}_{j+1} x^{p_{j+1}\omega_j} + \cdots + c_{j+1}\xi_{j+1}x^{\omega_{j+1}} + \text{l.o.t.}, \tag{19}$$

where $\tilde{c}_{j+1} := (c_j a_{j+1})^{p_{j+1}}$ and $c_{j+1} := p_{j+1} c_j^{p_{j+1}} a_{j+1}^{p_{j+1}-1}$. Note that $p_{j+1}\omega_j \in \frac{1}{p_1 \cdots p_j}$. Therefore, assertion 2 of lemma 5.6 implies that there exists $\alpha_0, \beta_{0,0}, \ldots, \beta_{0,j-1} \in \mathbb{Z}$ such that

1. $0 \leq \beta_{0,k} < p_{k+1}$ for $0 \leq k \leq j - 1$, and
2. $\alpha_0 + \sum_{k=0}^{j-1} \beta_{0,k}\omega_k = p_{j+1}\omega_j$.

Let

$$F_{j+1,0} := -\tilde{c}_{j+1} x^{\alpha_0} \prod_{k=0}^{j-1} \left(\frac{y_k}{c_k a_{k+1}}\right)^{\beta_{0,k}} \text{ and } H_{j+1,0} := y_j^{p_{j+1}} + F_{j+1,0}.$$

Now, (16) implies that for all k, $0 \leq k \leq j$,

$$\epsilon\left(f_k|_{y = \tilde{\psi}_{j+1}}\right) = r_{k+1} - r_{j+2}, \text{ and therefore,}$$

$$\epsilon\left(f_0|_{y = \tilde{\psi}_{j+1}}\right) > \cdots > \epsilon\left(f_{j-1}|_{y = \tilde{\psi}_{j+1}}\right) > \epsilon\left(f_j|_{y = \tilde{\psi}_{j+1}}\right) = r_{j+1} - r_{j+2} = \epsilon\left(f_j^{p_{j+1}}|_{y = \tilde{\psi}_{j+1}}\right) \tag{20}$$

Recall that $\pi_j : \mathbb{C}[x, x^{-1}, y_0, \ldots, y_j] \to \mathbb{C}[x,y]$ sends $x \mapsto x$ and $y_k \to f_k$ for $0 \le k \le j$. The above computations of $\epsilon(f_k)$'s imply that

$$\begin{gathered}
\pi(F_{j+1,0})|_{y=\tilde{\psi}_{j+1}} = -\tilde{c}_{j+1}x^{p_{j+1}\omega_j} + \cdots + c_{j+1,0}\xi_{j+1}x^{p_{j+1}\omega_j-\epsilon_0} + \text{l.o.t., where} \qquad (21)\\
c_{j+1,0} \in \mathbb{C} \text{ and } \epsilon_0 := \epsilon\left(\pi(F_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) \ge \min_{k=0}^{j-1} \epsilon\left(f_k|_{y=\tilde{\psi}_{j+1}}\right) > r_{j+1} - r_{j+2}.
\end{gathered}$$

Moreover, (18) implies that

$$c_\omega\left(\pi(F_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) \neq 0 \text{ and } \omega > p_{j+1}\omega_j - \epsilon_0 \quad \Rightarrow \quad \omega \in \langle 1, r_1, \ldots, r_{j+1}\rangle = \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}.$$

Since $p_{j+1}\omega_j - \epsilon_0 < p_{j+1}\omega_j - (r_{j+1} - r_{j+2}) = \omega_{j+1}$, it then follows from (19) and (21) that

$1_{(j+1,0)}$. $\tilde{\omega}_{j+1,0} := \deg_x\left(\pi(H_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) < p_{j+1}\omega_j$,

$2_{(j+1,0)}$. $\deg_{\mathrm{x}}^{\mathrm{inv}}\left(\pi(H_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) = \omega_{j+1}$, and $c_{\omega_{j+1}}\left(\pi(H_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) = c_{j+1}\xi_{j+1}$,

$3_{(j+1,0)}$. if $\omega > \omega_{j+1}$ and $c_\omega\left(\pi(H_{j+1,0})|_{y=\tilde{\psi}_{j+1}}\right) \neq 0$, then $\omega \in \langle 1, r_1, \ldots, r_{j+1}\rangle = \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}$.

If $\tilde{\omega}_{j+1,0} = \omega_{j+1}$, then set $F_{j+1} := H_{j+1,0}$. Otherwise $\tilde{\omega}_{j+1,0} > \omega_{j+1}$ and

$$\pi(H_{j+1,0})|_{y=\tilde{\psi}_{j+1}} = \tilde{c}_{j+1,0}x^{\tilde{\omega}_{j+1,0}} + \cdots + c_{j+1}\xi_{j+1}^{\omega_{j+1}} + \text{l.o.t.} \tag{22}$$

for some $\tilde{c}_{j+1,0} \neq 0 \in \mathbb{C}$. Then property $3_{(j+1,0)}$ implies that $\tilde{\omega}_{j+1,0} \in \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}$ and therefore assertion 2 of lemma 5.6 implies that there exists $\alpha_1, \beta_{1,0}, \ldots, \beta_{1,j} \in \mathbb{Z}$ such that

1. $0 \le \beta_{1,k} < p_{k+1}$ for $0 \le k \le j$, and
2. $\alpha_1 + \sum_{k=0}^{j} \beta_{1,k}\omega_k = \tilde{\omega}_{j+1,0}$.

Let

$$F_{j+1,1} := -\tilde{c}_{j+1,0}x^{\alpha_1}\prod_{k=0}^{j}\left(\frac{y_k}{c_k a_{k+1}}\right)^{\beta_{1,k}} \quad \text{and } H_{j+1,1} := H_{j+1,0} + F_{j+1,1}.$$

It then follows from (20) and (18) that

$$\begin{gathered}
\pi(F_{j+1,1})|_{y=\tilde{\psi}_{j+1}} = -\tilde{c}_{j+1,0}x^{\tilde{\omega}_{j+1,0}} + \cdots + c_{j+1,1}\xi_{j+1}x^{\tilde{\omega}_{j+1,0}-\epsilon_1} + \text{l.o.t., where} \qquad (23)\\
c_{j+1,1} \in \mathbb{C} \text{ and } \epsilon_1 := \epsilon\left(\pi(F_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) \ge \min_{k=0}^{j} \epsilon\left(f_k|_{y=\tilde{\psi}_{j+1}}\right) \ge r_{j+1} - r_{j+2}, \text{ and}\\
c_\omega\left(\pi(F_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) \neq 0 \text{ and } \omega > \tilde{\omega}_{j+1,0} - \epsilon_1 \quad \Rightarrow \quad \omega \in \langle 1, r_1, \ldots, r_{j+1}\rangle = \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}.
\end{gathered}$$

Since $\tilde{\omega}_{j+1,0} - \epsilon_1 < p_{j+1}\omega_j - \epsilon_1 \le p_{j+1}\omega_j - (r_{j+1} - r_{j+2}) = \omega_{j+1}$, it then follows from (22) and (23) that

$1_{(j+1,1)}$. $\tilde{\omega}_{j+1,1} := \deg_x\left(\pi(H_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) < \tilde{\omega}_{j+1,0}$,

$2_{(j+1,1)}$. $\deg_{\mathrm{x}}^{\mathrm{inv}}\left(\pi(H_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) = \omega_{j+1}$, and $c_{\omega_{j+1}}\left(\pi(H_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) = c_{j+1}\xi_{j+1}$,

$3_{(j+1,1)}$. if $\omega > \omega_{j+1}$ and $c_\omega\left(\pi(H_{j+1,1})|_{y=\tilde{\psi}_{j+1}}\right) \neq 0$, then $\omega \in \langle 1, r_1, \ldots, r_{j+1}\rangle = \frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}$.

If $\tilde{\omega}_{j+1,1} = \omega_{j+1}$, then set $F_{j+1} := H_{j+1,1}$. Otherwise $\tilde{\omega}_{j+1,1} > \omega_{j+1}$ and

$$\pi(H_{j+1,1})|_{y=\tilde{\psi}_{j+1}} = \tilde{c}_{j+1,1}x^{\tilde{\omega}_{j+1,0}} + \cdots + c_{j+1}\xi_{j+1}^{\omega_{j+1}} + \text{l.o.t.}$$

for some $\tilde{c}_{j+1,1} \neq 0 \in \mathbb{C}$, and it is straightforward to see that we may follow exactly the same procedure for constructing $H_{j+1,1}$ to construct $H_{j+1,2}$ from $H_{j+1,1}$, and then $H_{j+1,3}$ from $H_{j+1,2}$ and so on. More precisely, as long as $\deg_x\left(\pi(H_{j+1,i})|_{y=\tilde{\psi}_{j+1}}\right) > \omega_{j+1}$, following the above procedure we may construct $H_{j+1,i+1}$. It follows from the construction of $H_{j+1,i}$'s that $\{\deg_x\left(\pi(H_{j+1,i})|_{y=\tilde{\psi}_{j+1}}\right) : i \geq 0\}$ is a *decreasing* sequence of elements of $\frac{1}{p_1 \cdots p_{j+1}}\mathbb{Z}$ bounded below by ω_{j+1}, and therefore it is a *finite* sequence. Consequently, there exists $m_{j+1} \in \mathbb{Z}$ such that $\deg_x\left(\pi(H_{j+1,m_{j+1}})|_{y=\tilde{\psi}_{j+1}}\right) = \omega_{j+1}$. Let $F_{j+1} := H_{j+1,m_{j+1}}$. Assertions $1_{(j+1)}$, $2_{(j+1)}$, $3_{(j+1)}$, $4_{(j+1)}$ and $7_{(j+1)}$ are then immediate consequences of the construction of $H_{j+1,i}$'s.

It remains to prove assertions $5_{(j+1)}$ and $6_{(j+1)}$. The following lemma will be useful for this purpose.

Lemma 2.20. *If F is a η_j-homogeneous polynomial in A_j with $\deg_{y_i}(F) < p_{i+1}$ for all i, $0 \leq i \leq j$, then*

1. *every η_j-homogeneous form of F is a monomial, and*
2. *$\delta_{j+1}(F(x, f_0, \cdots, f_j)) = \eta_j(F)$.*

Proof. Assertion 1 of the claim is a direct consequence of assertion 2 of lemma 5.6. For assertion 1, note that $\delta_{j+1}(f_k) = \eta_j(y_k)$ for all k, $0 \leq k \leq j$ (due to (16)). It follows that $\delta_{j+1}(H(x, f_0, \ldots, f_j)) = \eta_j(H)$ for every monomial H in A_j. Let F be as in the statement of the lemma. Assertion 1 implies that $F = \sum F_i$ such that each F_i is a monomial with $\eta_j(F_1) > \eta_j(F_2) > \cdots$. It follows that $\delta_{j+1}(F_j(x, f_0, \ldots, f_j)) > \delta_{j+1}(F_{j+1}(x, f_0, \ldots, f_j)) > \cdots$ and consequently $\delta_{j+1}(F(x, f_0, \ldots, f_j)) = \delta_{j+1}(F_1(x, f_0, \ldots, f_j)) = \eta_j(F_1) = \eta_j(F)$, which is precisely the content of assertion 2. This completes the proof of the lemma. □

Let $F_{\psi_{j+1}}^{(j+1)}$ be as in corollary 5.4, i.e. $F_{\psi_{j+1}}^{(j+1)} \in A_j[x^{-1}]$ such that

$$F_{\psi_{j+1}}^{(j+1)}(x, f_0, \ldots, f_j) = \Psi_{j+1}^{(j+1)} = \Psi_{j+1}^*, \text{ and} \tag{P-1}$$

$$\deg_{y_k}(F_{\psi_{j+1}}^{(j+1)} - y_j^{p_{j+1}}) < p_{k+1} \text{ for all } k,\ 0 \leq k \leq j. \tag{P-2}$$

Claim 2.20.1. *We claim that F_{j+1} is precisely the sum of all monomial terms H of $F_{\psi_{j+1}}^{(j+1)}$ such that $\eta_j(F) > \omega_{j+1}$.*

Proof. Let $\tilde{F}_{j+1}$ be the sum of all monomial terms H of $F_{\psi_{j+1}}^{(j+1)}$ such that $\eta_j(F) > \omega_{j+1}$. Then $F_{\psi_{j+1}}^{(j+1)} = \tilde{F}_{j+1} + G_{j+1}$ such that $\eta_j(G_{j+1}) \leq \omega_{j+1}$. Now, (P-1), (13) and assertion $7_{(j+1)}$ imply that

$$\delta_{j+1}(\pi(\tilde{F}_{j+1} + G_{j+1} - F_{j+1})) = \delta_{j+1}(\Psi_{j+1}^* - f_{j+1}) \leq \max\{\delta_{j+1}(\Psi_{j+1}^*), \delta_{j+1}(f_{j+1})\} = \omega_{j+1}.$$

Since $\delta_{j+1}(\pi(G_{j+1})) \leq \eta_j(G_{j+1}) \leq \omega_{j+1}$, it follows that

$$\delta_{j+1}(\pi(\tilde{F}_{j+1} - F_{j+1})) \leq \omega_{j+1}.$$

Now note that $F := \tilde{F}_{j+1} - F_{j+1} = (\tilde{F}_{j+1} - y_j^{p_{j+1}}) - (F_{j+1} - y_j^{p_{j+1}})$. Therefore, due to (P-2) and assertion $3_{(j+1)}$, we see that F satisfies the hypothesis of lemma 2.20. If $F \neq 0$, then by definition of F_{j+1} and $\tilde{F}_{j+1}$, it follows that $\eta_j(F) > \omega_{j+1} \geq \delta_{j+1}(\pi(F))$, which contradicts assertion 2 of lemma 2.20. It follows that $F_{j+1} = \tilde{F}_{j+1}$, as required. □

Let G_{j+1} be as in the proof of claim 2.20.1. Then $F^{(j+1)}_{\psi_{j+1}} = F_{j+1} + G_{j+1}$, and therefore $\Psi^*_{j+1} = f_{j+1} + G_{j+1}(x, f_0, \ldots, f_j)$. Since $\eta_j(G_{j+1}) \leq \omega_{j+1}$, this proves assertion $6_{(j+1)}$.

It remains to prove assertion $5_{(j+1)}$. Note that Ψ^*_{j+1} is monic in y with $\deg_y(\Psi^*_{j+1}) = p_1p_2\cdots p_{j+1}$. On the other hand, inductive assumption $5_{(k)}$ implies that f_j is also monic in y with $\deg_y(f_j) = p_1p_2\cdots p_j$. Consequently, setting $\Phi := \Psi^*_{j+1} - f_j^{p_{j+1}}$, we see that $\deg_y(\Phi) < p_1p_2\cdots p_{j+1}$. Therefore, corollary 5.4 implies that there is a unique element $F_\Phi \in A_j[x^{-1}]$ such that $\Phi = F_\Phi(x, f_0, \ldots, f_j)$ and $\deg_{y_k}(F_\Phi) < p_{k+1}$ for all k, $0 \leq k \leq j$. But then assertion 2 of corollary 5.4 implies that $\Psi^*_{j+1} = y_j^{p_{j+1}} + F_\Phi$. Claim 2.20.1 then implies that every monomial term of $F_{j+1} - y_j^{p_{j+1}}$ is also a monomial term of F_Φ. The monomial terms of $F_{j+1} - y_j^{p_{j+1}}$ are precisely $F_{j+1,i}$, $0 \leq i \leq m_{j+1}$. Consequently, the last assertion of corollary 5.4 implies that for all i, $0 \leq i \leq m_{j+1}$,

$$\deg_y(\pi(F_{j+1,i})) \leq \deg_y(\Phi) < p_1p_2\cdots p_{j+1} = \deg_y(f_j^{p_{j+1}}).$$

Since $f_{j+1} = f_j^{p_{j+1}} + \sum_{i=0}^{m_{j+1}} \pi(F_{j+1,i})$, it follows that f_{j+1} is monic in y with $\deg_y(f_{j+1}) = p_1p_2\cdots p_{j+1}$. This proves assertion $5_{(j+1)}$ and completes the proof of claim 2.19.1. □

For $F \in B_l := \mathbb{C}\langle\langle x\rangle\rangle[y_0, \ldots, y_l]$ and $I \subseteq \mathbb{R}$, we write $F|_I$ for the sum of all monomial terms H of F such that $\eta(F) \in I$. Moreover, for $w \in \mathbb{R}$, we write $F|_{>w}$ for $F|_{(w,\infty)}$.

Lemma 2.21. *For each i, k such that $0 \leq i < k \leq l+1$, let $\omega_{i,k} := \omega_i + r_k - r_{i+1}$. Then*

1. *for each k, $1 \leq k \leq j$, F_k is precisely the sum of all monomial terms H of $F^{(k)}_{\psi_k}$ such that $\eta_{k-1}(H) > \omega_k = \omega_{k,k+1}$.*
2. *Let $\phi \in \mathbb{C}\langle\langle x\rangle\rangle$ such that $\phi \equiv_{r_{j'+1}} \psi$ with $j' \leq j+1$. Then for all i, k, $1 \leq i \leq k \leq j'$, $\eta_{i-1}(F^{(i)}_\phi - F^{(i)}_{\psi_k}) \leq \omega_{i,k+1}$.*

Proof. Let G_k, $1 \leq k \leq j$, be as in inductive assumption $6_{(k)}$. A direct application of corollary 5.4 (which is valid due to inductive assumptions $3_{(k)}$, $5_{(k)}$ and $6_{(k)}$) then shows that $F^{(k)}_{\psi_k} = F_k + G_k$ for all k, $1 \leq k \leq j$, which implies assertion 1.

For assertion 2, we prove by induction on i that

$$\eta_{i-1}(F^{(i)}_\phi - F^{(i)}_{\psi_k}) \leq \omega_{i,k+1}$$

for all pair (k, j') of integers and *all* $\phi \in \mathbb{C}\langle\langle x\rangle\rangle$ such that

$$1 \leq i \leq k \leq j' \leq j+1, \text{ and} \qquad (c_{i,k,j'})$$

$$\phi \equiv_{r_{j'+1}} \psi. \qquad (c_\phi)$$

Let $i = 1$. Pick $k, j' \in \mathbb{Z}$ and $\phi \in \mathbb{C}\langle\langle x\rangle\rangle$ such that conditions $(c_{i,k,j'})$ and (c_ϕ) are satisfied. Note that

$$\Phi^{(0)} := y - \phi(x) = y - \psi_k(x) - \hat{\phi}_0(x)$$

where $\hat{\phi}_0(x) := \phi(x) - \psi_k(x)$. Since $\phi \equiv_{r_{j'+1}} \psi$ and $j' \geq k$, it follows that $\deg(\hat{\phi}_0(x)) \leq r_{k+1}$. Moreover, the first k Puiseux pairs of ϕ are precisely (p_i, q_i) for $1 \leq i \leq k$. Therefore $p_1 \cdots p_k$ divides the polydromy order s of ϕ. Let μ be an s-th root of unity. Then

$$\begin{aligned}
\Phi^{(1)} &:= \prod_{m=1}^{p_1} (y - \mu^m \star_s \phi(x)) \\
&= \prod_{m=1}^{p_1} (y - \zeta_k^m \star_{p_1 \cdots p_k} \psi_k(x) - \mu^m \star_s \hat{\phi}_0(x)) \\
&= \Psi_k^{(1)} + G^{(1)}(x, y)
\end{aligned}$$

for some $G^{(1)} \in \mathbb{C}\langle\langle x\rangle\rangle[y_0]$ with $\eta_0(G^{(1)}) \leq (p_1 - 1)r_1 + r_{k+1} = \omega_{1,k+1}$. Note from the definition of $F_\phi^{(1)}$ that $F_\phi^{(1)} = \Phi^{(1)}(x, y_0)$ and $F_{\psi_k}^{(1)} = \Psi_k^{(1)}(x, y_0)$. It follows that $\eta_0(F_\phi^{(1)} - F_{\psi_k}^{(1)}) \leq \omega_{1,k+1}$ and the lemma is proved for $i = 1$.

Now assume that there exists i, $1 \leq i \leq j$, such that for all $k, j' \in \mathbb{Z}$ and $\phi \in \mathbb{C}\langle\langle x\rangle\rangle$ satisfying $(c_{i,k,j'})$ and (c_ϕ), $F_\phi^{(i)} = F_{\psi_k}^{(i)} + G_\phi^{k,i}$ for some $G_\phi^{k,i} \in \mathbb{C}\langle\langle x\rangle\rangle[y_0, \ldots, y_{i-1}]$ such that $\eta_{i-1}(G_\phi^{k,i}) \leq \omega_{i,k+1}$. Fix $\tilde{k}$, $\tilde{j}$ and $\tilde{\phi}$ such that

$$i + 1 \leq \tilde{k} \leq \tilde{j} \leq j + 1, \text{ and} \qquad (c_{i+1,\tilde{k},\tilde{j}})$$

$$\tilde{\phi} \equiv_{r_{\tilde{j}+1}} \psi. \qquad (c_{\tilde{\phi}})$$

Note that $(c_{i,k,j'})$ and (c_ϕ) are satisfied with $k = i$, $j' = \tilde{k}$ and $\phi = \psi_{\tilde{k}}$. Therefore, by induction hypothesis,

$$F_{\psi_{\tilde{k}}}^{(i)} = F_{\psi_i}^{(i)} + G_{\psi_{\tilde{k}}}^{i,i}, \text{ where } \eta_{i-1}\left(G_{\psi_{\tilde{k}}}^{i,i}\right) \leq \omega_{i,i+1} = \omega_i.$$

Assertion 1 then implies that

$$F_{\psi_{\tilde{k}}}^{(i)} = F_i + G_{\tilde{k},i}, \text{ where } \eta_{i-1}\left(G_{\tilde{k},i}\right) \leq \omega_i. \qquad (c_{\tilde{k},i})$$

Now applying the induction hypothesis with $k = \tilde{k}$, $j' = \tilde{j}$ and $\phi = \tilde{\phi}$, we see that

$$F_{\tilde{\phi}}^{(i)} = F_{\psi_{\tilde{k}}}^{(i)} + G_{\tilde{k},i}^{\tilde{\phi}} \text{ (where } \eta_{i-1}\left(G_{\tilde{k},i}^{\tilde{\phi}}\right) \leq \omega_{i,\tilde{k}+1}).$$

Since $F_{\psi_{\tilde{k}}}^{(i)}|_{>\omega_i} = F_i$, applying lemma 2.22 with $k = i$, $\phi = \psi_{\tilde{k}}$ and $w = \omega_{i,\tilde{k}+1}$ shows that

$$F_{\tilde{\phi}}^{(i+1)} = F_\phi^{(i+1)} + G$$

with $\eta(G) \leq (p_{i+1} - 1)\omega_i + \omega_{i,\tilde{k}+1} = \omega_{i+1,\tilde{k}+1}$. This completes the proof of the lemma by induction. □

Lemma 2.22. *Let* $1 \le k \le j$ *and* $\phi \in \mathbb{C}\langle\langle x \rangle\rangle$ *such that* $\phi \equiv_{r_{k+2}} \psi$ *and* $F_\phi^{(k)}|_{>\omega_k} = F_k$*. Then for each* $w < \omega_k \in \mathbb{R}$, $F_\phi^{(k+1)}|_{>(p_{k+1}-1)\omega_k + w}$ *is uniquely determined by* $F_\phi^{(k)}|_{>w}$*.*

Proof. Let $G := F_\phi^{(k)}|_{(w,\omega_k]}$ and $\hat{G} := F_\phi^{(k)}|_{(-\infty,w]}$. Note that $\eta(F_k) > \omega_k$ (due to inductive assumption $4_{(k)}$). It follows that $F_\phi^{(k)} = F_k + G + \hat{G}$.

Let s be the polydromy order of ϕ and μ be an s-th root of unity. Then

$$\begin{aligned}
\Phi^{(k+1)} &= \prod_{n=0}^{p_1 \cdots p_{k+1}-1} (y - \mu^n \star_s \phi(x)) = \prod_{n=0}^{p_1 \cdots p_k - 1} \prod_{m=0}^{p_{k+1}-1} \left(y - \mu^{mp_1 \cdots p_k + n} \star_s \phi(x)\right) \\
&= \prod_{m=0}^{p_{k+1}-1} \mu^{mp_1 \cdots p_k} \star_s \left(\prod_{n=0}^{p_1 \cdots p_k - 1} (y - \mu^n \star_s \phi(x)) \right) = \prod_{m=0}^{p_{k+1}-1} \mu^{mp_1 \cdots p_{k+1}} \star_s \Phi^{(k)} \\
&= \tilde{H}^\phi(x, f_0, \ldots, f_{k-1}), \text{ where} \\
\tilde{H}^\phi &:= \prod_{m=0}^{p_{k+1}-1} \mu^{mp_1 \cdots p_k} \star_s F_\phi^{(k)} \in \mathbb{C}\langle\langle x \rangle\rangle[y_0, \ldots, y_{k-1}] =: B_{k-1}.
\end{aligned}$$

Note that

$$\begin{aligned}
\tilde{H}^\phi &= \prod_{m=0}^{p_{k+1}-1} \mu^{mp_1 \cdots p_k} \star_s (F_k + G + \hat{G}) = \prod_{m=0}^{p_{k+1}-1} (F_k + \mu^{mp_1 \cdots p_k} \star_s G + \mu^{mp_1 \cdots p_k} \star_s \hat{G}) \\
&= \prod_{m=0}^{p_{k+1}-1} (F_k + \lambda^{mp_1 \cdots p_k} \star_t G + \mu^{mp_1 \cdots p_k} \star_s \hat{G}),
\end{aligned}$$

where t is the polydromy order of G and λ is a t-th root of unity. Let

$$H_0^\phi := \prod_{m=0}^{p_{k+1}-1} \left(y_k + \lambda^{mp_1 \cdots p_k} \star_t G + \mu^{mp_1 \cdots p_k} \star_s \hat{G}\right) \in \mathbb{C}\langle\langle x \rangle\rangle[y_0, \ldots, y_k] =: B_k.$$

Then $H_0^\phi(x, f_0, \ldots, f_k) = \tilde{H}^\phi(x, f_0, \ldots, f_{k-1}) = \Phi^{(k+1)}$ and $\deg_{y_k}(H_0^\phi - y_k^{p_{k+1}}) < p_{k+1}$. Let $\prec_k$ be the monomial ordering on B_k introduced in the set-up for induction. Recall that $\prec_k$ refines the partial ordering on monomials induced by η_k. Since $\eta_k(G + \hat{G}) \le \omega_k = \eta_k(y_k)$ and since y_k does *not* appear in any of the monomials in G or $\hat{G}$, it follows from the definition of $\prec_k$ that the maximum (w.r.t. $\prec_k$) of all monomials which appear in H_0^ϕ is $y_k^{p_{k+1}}$, or equivalently, the following property holds for $n = 0$ (where for every $\alpha \in \mathbb{Z}$, $\beta := (\beta_0, \ldots, \beta_k) \in \mathbb{Z}^{k+1}$, and $\tau \in B_k$, we write $c_{\alpha,\beta}(\tau)$ for the coefficient of $x^\alpha y_0^{\beta_0} \cdots y_k^{\beta_k}$ in τ):

$$y_k^{p_{k+1}} \succ_k x^\alpha y_0^{\beta_0} \cdots y_k^{\beta_k} \text{ for all } \alpha \in \mathbb{Z} \text{ and } \beta \in \mathbb{Z}^{k+1} \text{ such that } c_{\alpha,\beta}(H_n^\phi - y_k^{p_{k+1}}) \neq 0 \qquad (P_n)$$

Starting with H_0^ϕ, we now inductively construct a sequence $\{H_n^\phi : n \ge 0\} \subseteq B_k$ such that each H_n^ϕ satisfies (P_n).

So assume H_n^ϕ is given and H_n^ϕ satisfies (P_n). Write $H_n^\phi - y_k^{p_{k+1}} = \sum c_{n,\beta}^\phi(x) y_0^{\beta_0} \cdots y_k^{\beta_k}$, i.e. $c_{n,\beta}^\phi(x) := \sum_\alpha \left(c_{n,\alpha,\beta}^\phi(H_n^\phi - y_k^{p_{k+1}})\right) x^\alpha$. Let

$$S_n^\phi := \{\beta \in \mathbb{Z}^{k+1} : c_{n,\beta}^\phi(x) \not\equiv 0 \text{ and } \beta_i \ge p_{i+1} \text{ for some } i,\ 0 \le i \le k\}.$$

Set $H^{\phi}_{n+1} := H^{\phi}_n$ if $S^{\phi}_n = \emptyset$. Otherwise, let $\beta^{\phi,n} := \max\{\beta : \beta \in S^{\phi}_n\}$, where the maximum is taken with respect to $\prec_k$. Let $\hat{i}_n$ be the 'right-most' coordinate of $\beta^{\phi,n}$ such that $\beta^{\phi,n}_{\hat{i}_n} \geq p_{\hat{i}_n+1}$. We claim that $\hat{i}_n < k$. Indeed, since δ is positive on non-constant polynomials in $\mathbb{C}[x,y]$, inductive assumptions $5_{(k)}$ and $7_{(k)}$ imply that $\eta_k(y_i) = \delta(f_i) > 0$ for all i, $0 \leq i \leq k$. Then (P_n) implies that $y_0^{\beta^{\phi,n}_0} \cdots y_k^{\beta^{\phi,n}_k} \prec_k y_k^{p_{k+1}}$ and therefore $\beta^{\phi,n}_k < p_{k+1}$. Consequently, $\hat{i}_n < k$, as claimed. In particular, $F_{\hat{i}_n+1}$ is defined and satisfies inductive assumptions $0_{(k)}$ to $9_{(k)}$. Let

$$M_{\beta^{\phi,n}} := y_0^{\beta^{\phi,n}_0} \cdots y_{\hat{i}_n-1}^{\beta^{\phi,n}_{\hat{i}_n-1}} y_{\hat{i}_n}^{\beta^{\phi,n}_{\hat{i}_n}-p_{\hat{i}_n+1}} y_{\hat{i}_n+1}^{\beta^{\phi,n}_{\hat{i}_n+1}} \cdots y_k^{\beta^{\phi,n}_k} \left(y_{\hat{i}_n+1} - \left(F_{\hat{i}_n+1} - y_{\hat{i}_n}^{p_{\hat{i}_n+1}} \right) \right), \text{and}$$

$$H^{\phi}_{n+1} := y_k^{p_{k+1}} + \sum_{\beta \neq \beta^{\phi,n}} c^{\phi}_{n,\beta}(x) y_0^{\beta_0} \cdots y_k^{\beta_k} + c^{\phi}_{n,\beta^{\phi,n}}(x) M_{\beta^{\phi,n}}.$$

Inductive assumption $4_{(k)}$ implies that $y_{\hat{i}_n}^{p_{\hat{i}_n+1}} \succ_k M$ for every monomial M in $F_{\hat{i}_n+1} - y_{\hat{i}_n}^{p_{\hat{i}_n+1}}$. Morevoer, $y_{\hat{i}_n}^{p_{\hat{i}_n+1}} \succ_k y_{\hat{i}_n+1}$, since $\eta(y_{\hat{i}_n+1}) = \omega_{\hat{i}_n+1} < p_{\hat{i}_n+1}\omega_{\hat{i}_n} = \eta(y_{\hat{i}_n}^{p_{\hat{i}_n+1}})$ (according to (35)). Pick $(\beta_0, \cdots, \beta_k)$ such that coefficient of $y_0^{\beta_0} \cdots y_k^{\beta_k}$ in $M_{\beta^{\phi,n}}$ is non-zero. Then it follows that $y_0^{\beta_0} \cdots y_k^{\beta_k} \prec_k y_0^{\beta^{\phi,n}_0} \cdots y_k^{\beta^{\phi,n}_k}$, and therefore, for all α such that the coefficient of x^{α} in $c^{\phi}_{n,\beta^{\phi,n}}(x)$ is non-zero,

$$y_k^{p_{k+1}} \succ_k x^{\alpha} y_0^{\beta^{\phi,n}_0} \cdots y_k^{\beta^{\phi,n}_k} \text{ (due to } (P_n)) \succ_k x^{\alpha} y_0^{\beta_0} \cdots y_k^{\beta_k}.$$

In particular, (P_{n+1}) is satisfied.

We claim that the sequence of H^{ϕ}_n's stabilize at a finite step. Indeed, for all $n \geq 0$, if $H^{\phi}_{n+1} \neq H^{\phi}_n$, then the construction of H^{ϕ}_{n+1} shows that $\beta^{\phi,n+1} \prec_k \beta^{\phi,n}$. Since $\prec_k$ induces a well-ordering on $\mathbb{Z}^{k+1}_{\geq 0}$ (see the remark preceding inductive assumption $9_{(k)}$), it follows that the sequence $\{\beta^{\phi,n} : n \geq 0\}$ has to stop at a finite step, and consequently there exists $n_0 \geq 0$ such that $H^{\phi}_n = H^{\phi}_{n_0}$ for all $n \geq n_0$. Then $S^{\phi}_{n_0} = \emptyset$ and therefore $\deg_{y_i}(H^{\phi}_{n_0} - y_k^{p_{k+1}}) < p_{i+1}$ for all i, $0 \leq i \leq k$. Moreover, note that for all $n \geq 0$, $H^{\phi}_{n+1}(x, f_0, \ldots, f_k) = H^{\phi}_n(x, f_0, \ldots, f_k)$, and therefore, $H^{\phi}_{n_0}(x, f_0, \ldots, f_k) = H^{\phi}_0(x, f_0, \ldots, f_k) = \Phi^{(k+1)}$. It follows that $H^{\phi}_{n_0} = F^{(k+1)}_{\phi}$.

Now pick $\tilde{\phi} \in \mathbb{C}\langle\langle x \rangle\rangle$ such that $\tilde{\phi} \equiv_{r_{k+2}} \psi$ and $F^{(k)}_{\tilde{\phi}}|_{>w} = F^{(k)}_{\phi}|_{>w} = F_k + G$. Let $w^* := (p_k - 1)\omega_{k+1} + w$. We have to show that $F^{(k+1)}_{\phi}|_{>w^*} = F^{(k+1)}_{\tilde{\phi}}|_{>w^*}$.

By our assumption, $F^{(k)}_{\tilde{\phi}} = F_k + G + \tilde{G}$ for some $\tilde{G} \in B_{k-1}$ such that $\eta_{k-1}(\tilde{G}) \leq w$. Therefore, if $\tilde{s}$ is the polydromy order of $\tilde{\phi}$ and $\tilde{\mu}$ is a $\tilde{s}$-th root of unity, then

$$H^{\tilde{\phi}}_0 := \prod_{m=0}^{p_{k+1}-1} \left(y_k + \lambda^{mp_1 \cdots p_k} \star_t G + \tilde{\mu}^{mp_1 \cdots p_k} \star_{\tilde{s}} \tilde{G} \right).$$

Note that all monomial terms M in $H^{\tilde{\phi}}_0$ with $\eta(M) > w^*$ are also monomial terms of $\prod_{m=0}^{p_{k+1}-1} (y_k + \lambda^{mp_1 \cdots p_k} \star_t G)$. It follows that $H^{\phi}_0|_{>w^*} = H^{\tilde{\phi}}_0|_{>w^*}$. Now we show by induction on n that $H^{\phi}_n|_{>w^*} = H^{\tilde{\phi}}_n|_{>w^*}$ for all $n \geq 0$. Indeed, we have just seen that the claim is true for $n = 0$. So assume it holds for $n \geq 0$. Let

$$M^{\phi} := \begin{cases} 0 & \text{if } S^{\phi}_n = \emptyset, \\ \sum_{\eta(x^{\alpha} y^{\beta^{\phi,n}}) > w^*} c_{\alpha,\beta^{\phi,n}}(\phi) x^{\alpha} y^{\beta^{\phi,n}} & \text{otherwise,} \end{cases}$$

where we wrote $y^{\beta^{\phi,n}}$ as a shortcut for $y_0^{\beta_0^{\phi,n}} \cdots y_k^{\beta_k^{\phi,n}}$. Recall that all the monomials in $\left(H^{\phi}_{n+1} - H^{\phi}_n\right)\Big|_{>w^*}$ arises from substitution of a polynomial $M_{\beta^{\phi,n}}$ for $y^{\beta^{\phi,n}}$ in M^{ϕ}. Now, note that the induction hypothesis implies that if $M^{\phi} \neq 0$, then $M^{\phi} = M^{\tilde{\phi}}$, and therefore $M_{\beta^{\phi,n}} = M_{\beta^{\tilde{\phi},n}}$. It follows that

$$\left(H^{\phi}_{n+1} - H^{\phi}_n\right)\Big|_{>w^*} = \left(H^{\tilde{\phi}}_{n+1} - H^{\tilde{\phi}}_n\right)\Big|_{>w^*},$$

and therefore $H^{\phi}_{n+1}\Big|_{>w^*} = H^{\tilde{\phi}}_{n+1}\Big|_{>w^*}$, which completes the proof of the claim by induction.

Since H^{ϕ}_n's (resp. $H^{\tilde{\phi}}_n$'s) eventually stabilize to $F^{(k+1)}_{\phi}$ (resp. $F^{(k+1)}_{\tilde{\phi}}$), it follows that $F^{(k+1)}_{\phi}|_{>w^*} = F^{(k+1)}_{\tilde{\phi}}|_{>w^*}$, as required. □

Claim 2.22.1. *Let* $F_{j+1} = y_j^{p_{j+1}} + \sum_{i=0}^{m_{j+1}} F_{j+1,i}$ *be as in claim 2.19.1. Assume there exists* i, $0 \leq i \leq m_{j+1}$, *such that* $\deg_x(F_{j+1,i})$ *is negative. If* $\phi_1, \ldots, \phi_m$ *are degree-wise Puiseux series in* x *such that* $\phi_j \equiv_{r_{j+2}} \psi$ *for each* j, $1 \leq n \leq m$, *then* $\prod_{n=1}^{m} \Phi^*_n \notin \mathbb{C}[x,y]$.

Proof. Let $\phi_1, \ldots, \phi_m$ be as in the statement of the lemma. Assume to the contrary of the claim that $\prod_{n=1}^{m} \Phi^*_n \in \mathbb{C}[x,y]$. Pick n, $1 \leq n \leq m$. Let s_n denote the polydromy order of ϕ_n and μ_n be a s_n-th root of unity. Since $\phi_n \equiv_{r_{j+2}} \psi$, the first $j+1$ Puiseux pairs of ϕ_n are (p_i, q_i), $1 \leq i \leq j+1$. In particular, $p_1 p_2 \cdots p_{j+1}$ divides s_n. Now, applying assertion 2 of lemma 2.21 with $\phi = \phi_n$ and $i = k = j' = j+1$ implies that $\eta_j(F^{(j+1)}_{\phi_n} - F^{(j+1)}_{\psi_{j+1}}) \leq \omega_{j+1,j+2} = \omega_{j+1}$. Claim 2.20.1 then shows that

$$F^{(j+1)}_{\phi_n} = F_{j+1} + \tilde{G}_n,$$

where $\tilde{G}_n \in B_j := \mathbb{C}\langle\langle x \rangle\rangle[y_0, \ldots, y_j]$ and $\eta_j(\tilde{G}_n) \leq \omega_{j+1}$. Let $t_n := \frac{s_n}{p_1 p_2 \cdots p_{j+1}}$. Then t_n is a positive integer and

$$\begin{aligned} \Phi^*_n &:= \prod_{i=0}^{s_n - 1} \mu_n^i \star_{s_n} (y - \phi_n(x)) = \prod_{i=0}^{t_n - 1} \prod_{k=0}^{p_1 p_2 \cdots p_{j+1} - 1} \mu_n^{i p_1 p_2 \cdots p_{j+1} + k} \star_{s_n} (y - \phi_n(x)) \\ &= \prod_{i=0}^{t_n - 1} \mu_n^{i p_1 p_2 \cdots p_{j+1}} \star_{s_n} \left(\prod_{k=0}^{p_1 p_2 \cdots p_{j+1} - 1} \mu_n^k \star_{s_n} (y - \phi_n(x)) \right) = \prod_{i=0}^{t_n - 1} \mu_n^{i p_1 p_2 \cdots p_{j+1}} \star_{s_n} \Phi^{(j+1)}_n \\ &= G^*_n(x, f_0, \ldots, f_j), \text{ where } G^*_n \in B_j \text{ is defined as} \\ G^*_n &:= \prod_{i=0}^{t_n - 1} \mu_n^{i p_1 p_2 \cdots p_{j+1}} \star_{s_n} F^{(j+1)}_{\phi_n} = \prod_{i=0}^{t_n - 1} (F_{j+1} + \mu_n^{i p_1 p_2 \cdots p_{j+1}} \star_{s_n} \tilde{G}_n). \end{aligned}$$

Let k_0 be the smallest integer such that $\deg_x(F_{j+1,k_0}) < 0$ and $w_0 := \eta_j(F_{j+1,k_0})$. Then $w_0 > \omega_{j+1}$ and therefore

$$G^*_n = \prod_{i=0}^{t_n - 1} (y_j^{p_{j+1}} + F_{j+1,0} + \cdots + F_{j+1,k_0} + \tilde{H}_{n,i}),$$

for some $\tilde{H}_{n,i} \in B_j$ with $\eta_j(\tilde{H}_{n,i}) < w_0$. Let $M := \sum_{n=1}^m t_n$ and for each i, $1 \leq i \leq M$, let $H_i := \tilde{H}_{n,k}$, where $i = t_1 + \cdots + t_{n-1} + k + 1$ and $0 \leq k < t_n$. Then $f = G^*(x, f_0, \ldots, f_j)$, where

$$G^* := \prod_{n=1}^{m} G_n^* = \prod_{i=1}^{M} (y_j^{p_{j+1}} + F_{j+1,0} + \cdots + F_{j+1,k_0} + H_i)$$

Let

$$G := \begin{cases} G^* - y_j^{Mp_{j+1}} & \text{if } k_0 = 0, \\ G^* - (y_j^{p_{j+1}} + F_{j+1,0} + \cdots + F_{j+1,k_0-1})^M & \text{otherwise.} \end{cases}$$

Then

$$\mathfrak{L}_{\eta_j}(G) = \begin{cases} F_{j+1,0} & \text{if } k_0 = 0 \text{ and } M = 1, \\ M y_j^{(M-1)p_{j+1}} F_{j+1,0} + \sum_{i=2}^{M} \binom{M}{i} y_j^{(M-i)p_{j+1}} F_{j+1,0}^{i} & \text{if } k_0 = 0, \text{and } M > 1, \\ M(y_j^{p_{j+1}} + F_{j+1,0})^{M-1} F_{j+1,k_0} & \text{otherwise.} \end{cases} \tag{24}$$

Recall that by our assumption $\pi_j(G^*) = \prod_{n=1}^m \Phi_n^* \in \mathbb{C}[x, y]$. Therefore $f := \pi_j(G)$ is also a polynomial in x and y. According to inductive assumption $8_{(k)}$ (applied with $k = j$), δ_j is induced by η_j via π_j. Therefore there is a polynomial $F \in A_j$ such that $\pi_j(F) = f$ and $\eta_j(F) = \delta_j(f) \leq \eta_j(G)$ (where the last inequality holds because $\delta_j(\pi_j(H)) \leq \eta_j(H)$ for all $H \in B_j$).

Claim 2.22.2. *$\eta_j(F) = \eta_j(G)$.*

Proof. Let $\prec_j$ be the monomial ordering in A_j defined in the set-up for induction. Let $b \in \mathbb{C}$ and $\alpha, \beta_0, \ldots, \beta_j \in \mathbb{Z}$ such that $F_{j+1,k_0} = bx^{-\alpha} y_0^{\beta_0} \cdots y_j^{\beta_j}$. Then $\alpha > 0$ due to our choice of k_0 and $0 < \beta_i < p_{i+1}$ for $0 \leq i \leq j$ due to assertion $3_{(j+1)}$ of claim 2.19.1. Assertion $2_{(j+1)}$ of claim 2.19.1 implies that if $k_0 = 0$ then $\beta_j = 0$. Therefore, if k is the smallest positive integer such that $x^k \mathfrak{L}_{\eta_j}(G) \in \mathbb{C}[x, y]$, then (24) implies that

$$\mathrm{LT}_{\prec_j}(\mathfrak{L}_{\eta_j}(x^k G)) = \begin{cases} b y_0^{\beta_0} \cdots y_{j-1}^{\beta_{j-1}} & \text{if } k_0 = 0 \text{ and } M = 1, \\ M b x^{(M-1)\alpha} y_0^{\beta_0} \cdots y_{j-1}^{\beta_{j-1}} y_j^{(M-1)p_{j+1}} & \text{if } k_0 = 0, \text{and } M > 1, \\ M b y_0^{\beta_0} \cdots y_{j-1}^{\beta_{j-1}} y_j^{(M-1)p_{j+1} + \beta_j} & \text{otherwise.} \end{cases} \tag{24'}$$

Let J_j be the ideal in A_j generated by all weighted homogeneous (w.r.t. η_j) polynomials f such that $\delta_j(f(x, f_0, \ldots, f_j)) < \eta_j(f)$. According to inductive assumption $9_{(k)}$ (applied with $k = j$), $\mathcal{B}_j := (y_{j-1}^{p_j} + F_{j,0}, y_{j-2}^{p_{j-1}} + F_{j-1,0}, \ldots, y_0^{p_1} + F_{1,0})$ is a Gröbner basis of J_j with respect to $\prec_j$. Assume to the contrary of the claim that $\eta_j(F) < \eta_j(G)$. Then

$$\delta_j(\pi_j(\mathfrak{L}_{\eta_j}(G))) = \delta_j(\pi_j(F) - \pi_j(G - \mathfrak{L}_{\eta_j}(G))) < \eta_j(G).$$

Hence, $x^k \mathfrak{L}_{\eta_j}(G) \in J_j$ and therefore there exists i, $1 \leq i \leq j$, such that $y_{i-1}^{p_i} = \mathrm{LT}_{\prec_j}(y_{i-1}^{p_i} + F_{i,0})$ divides $\mathrm{LT}_{\prec_j}(x^k \mathfrak{L}_{\eta_j}(G)) = \mathrm{LT}_{\prec_j}(\mathfrak{L}_{\eta_j}(x^k G))$. On the other hand, (24′) implies that $y_{i-1}^{p_i}$ does *not* divide $\mathrm{LT}_{\prec_j}(\mathfrak{L}_{\eta_j}(x^k G))$ for all i, $1 \leq i \leq j$. This contradiction proves the claim. ☐

Claim 2.22.2 implies that $\eta_j(F) = \eta_j(G)$ and therefore $H := \mathfrak{L}_{\eta_j}(x^k G) - \mathfrak{L}_{\eta_j}(x^k F) \in J_j$, where n is as in claim 2.22.2. Let

$$H_1 := \mathrm{LT}_{\prec_j}(\mathfrak{L}_{\eta_j}(x^k G)) \text{ and } H_2 := \mathrm{LT}_{\prec_j}(\mathfrak{L}_{\eta_j}(x^k F)).$$

Now, note that by our choice of F, $\mathfrak{L}_{\eta_j}(x^kF) \notin J_j$. Therefore, 'dividing out' F by $\mathcal{B}_j$ (which does not change $\eta_j(F)$), we may assume that $y_{i-1}^{p_i}$ does *not* divide H_2 for all i, $1 \leq i \leq j$. On the other hand, since $F \in \mathbb{C}[x,y]$, it follows that

$$\deg_x(H_2) \geq k > \deg_x(H_1).$$

In particular, $H_1 \neq H_2$ and therefore,

$$\mathrm{LT}_{\prec_j}(H) = \max\{H_1, -H_2\}.$$

It then follows from $(24')$ that $y_{i-1}^{p_i}$ does *not* divide $\mathrm{LT}_{\prec_j}(H)$ for all i, $1 \leq i \leq j$. Since $\mathcal{B}_j$ is a Gröbner basis of J_j with respect to $\prec_j$, it follows as in the proof of claim 2.22.2 that $H \notin J_j$. This gives rise to a contradiction and proves the claim. ☐

Now assume $F_{j+1} \in A_j$, i.e. $\deg_x(F_{j+1,i}) \geq 0$ for all i, $0 \leq i \leq m_{j+1}$. Then $f_{j+1} \in \mathbb{C}[x,y]$. Since δ is positive on non-constant polynomials in x and y, assertions $5_{(j+1)}$ and $7_{(j+1)}$ imply that $\omega_{j+1} > 0$. Since $\omega_k = \delta(f_k) > 0$ for all k, $0 \leq k \leq j$, it follows that $\prec_{j+1}$ induces an monomial ordering on A_{j+1}. We claim:

Claim 2.22.3.

$8_{(j+1)}$. *δ_{j+1} is precisely the degree-like function on $\mathbb{C}[x,y]$ induced by η_{j+1} via π_{j+1}, where $\pi_{j+1} : A_{j+1} \to \mathbb{C}[x,y]$ is the $\mathbb{C}$-algebra homomorphism that sends $x \mapsto x$ and $y_i \mapsto f_i$ for* $0 \leq i \leq j+1$.

$9_{(j+1)}$. *Let J_{j+1} be the ideal in A_{j+1} generated by all weighted homogeneous (w.r.t. η_{j+1}) polynomials f such that $\delta_{j+1}(f(x, f_0, \ldots, f_{j+1})) < \eta_{j+1}(f)$. Then $\mathcal{B}_{j+1} := (y_j^{p_{j+1}} + F_{j+1,0}, y_{j-1}^{p_j} + F_{j,0}, \ldots, y_0^{p_1} + F_{1,0})$ is a Gröbner basis of J_{j+1} with respect to $\prec_{j+1}$.*

$10_{(j+1)}$. *There exists an isomorphism $\chi_{j+1} : A_{j+1}/J_{j+1} \cong R_{j+1} \subseteq \mathbb{C}[x_{j+1}, y_{j+1}] =: S_{j+1}$ such that χ_{j+1} sends $x \mapsto x_{j+1}^{p_1 \cdots p_{j+1}}$, $y_{j+1} \mapsto y_{j+1}$, and $y_i \mapsto b_i x_{j+1}^{p_1 \cdots p_{j+1} \omega_i}$ for some $b_i \neq 0 \in \mathbb{C}$ for each i, $0 \leq i \leq j$. Consider A_{j+1} and S_{j+1} as graded rings, the grading on A_{j+1} being induced by η_{j+1} and the grading on S_{j+1} being induced by the weighted degree with weights $\frac{1}{p_1 \cdots p_{j+1}}$ for x_{j+1} and ω_{j+1} for y_{j+1}. Then R_{j+1} is a graded subring of S_{j+1} and χ_{j+1} is an isomorphism of graded rings.*

Proof. Let J_j be the ideal in A_j as in inductive assumption $10_{(k)}$ (for $k = j$). Applying $10_{(k)}$ with $k = j$, we see that there exists an isomorphism $\chi_j : A_j/J_j \cong R_j \subseteq \mathbb{C}[x_j, y_j] =: S_j$ such that χ_j sends $x \mapsto x_j^{p_1 \cdots p_j}$, $y_j \mapsto y_j$, and $y_i \mapsto b_i x_j^{p_1 \cdots p_j \omega_i}$ for some $b_i \in \mathbb{C}$ for each i, $0 \leq i \leq j-1$. Let $\tilde{K}_j$ be the ideal of A_j generated by J_j and $H_{j+1} := y_j^{p_{j+1}} + F_{j+1,0}$. At first we show that $\tilde{K}_j$ is a prime ideal of A_j and determine the quotient $A_j/\tilde{K}_j$.

Assertion $2_{(j+1)}$ implies that there exists $b \in \mathbb{C}$ and $\alpha, \beta_0, \ldots, \beta_{j-1} \in \mathbb{Z}_{\geq 0}$ such that $F_{j+1,0} = bx^\alpha y_0^{\beta_0} \cdots y_{j-1}^{\beta_{j-1}}$. Moreover, assertion $4_{(j+1)}$ implies that $\alpha + \sum_{i=0}^{j-1} \omega_i \beta_i = p_{j+1}\omega_j$. It follows that χ_j maps (the equivalence class of) H_{j+1} to $y_j^{p_{j+1}} + b_j x_j^{p_1 \cdots p_{j+1} \omega_j}$, where $b_j := bb_0^{\beta_0} \cdots b_{j-1}^{\beta_{j-1}}$. Now, by definition,

$$\omega_j = \frac{s_j}{p_1 \cdots p_j} + \frac{q_{j+1}}{p_1 \cdots p_{j+1}}$$

for some $s_j \in \mathbb{Z}$. It follows that $\gcd(p_{j+1}, p_j \cdots p_{j+1}\omega_j) = \gcd(p_{j+1}, q_{j+1}) = 1$. Consequently, $\chi_j(H_{j+1})$ is an irreducible element of S_j and the ideal L_j in S_j generated by $\chi_j(H_{j+1})$ is prime. We claim that $\chi_j(\tilde{K}_j) = L_j \cap R_j$. Indeed, $\chi_j(\tilde{K}_j)$ is precisely the ideal generated by $\chi_j(H_{j+1})$ in R_j and therefore $\chi_j(\tilde{K}_j) \subseteq L_j \cap R_j$. To see the inclusion in the other direction, recall (from inductive assumption $10_{(k)}$ for $k = j$) that $\chi_j : A_j/J_j \hookrightarrow S_j$ is a homomorphism of graded rings, where the grading on A_j is induced by η_j and S_j is graded by the weighted degree with weights $\frac{1}{p_1 \cdots p_j}$ for x_j and ω_j for y_j. Note that $\chi_j(H_{j+1})$ is a homogeneous element of S_j, so that L_j is a homogeneous ideal of S_j. Since R_j is a graded subring of S_j, it suffices to show that every homogeneous element of $L_j \cap R_j$ belongs to $\chi_j(\tilde{K}_j)$ Let h be a non-zero homogeneous element of $L_j \cap R_j$. Then

$$h = a' x_j^{\alpha'} y_j^{\beta'} (y_j^{p_{j+1}} + b_j x_j^{p_1 \cdots p_{j+1}\omega_j}) \prod_{i=1}^{k} (y_j^{p_{j+1}} - b'_i x_j^{p_1 \cdots p_{j+1}\omega_j})$$

for non-negative integers α', β' and non-zero complex numbers $a', b'_1, \ldots, b'_k$. Note that the coefficient of $x_j^{\alpha'} y_j^{(k+1)p_{j+1}+\beta'}$ in h is non-zero. Since $h \in R_j$, it follows that $x_j^{\alpha'} y_j^{(k+1)p_{j+1}+\beta'} \in R_j$. Now, R_j is by definition the image of χ_j, so that

$$R_j = \mathbb{C}[x_j^{p_1 \cdots p_j}, x_j^{p_1 \cdots p_j \omega_0}, \cdots, x_j^{p_1 \cdots p_j \omega_{j-1}}, y_j].$$

Since $x_j^{\alpha'} y_j^{(k+1)p_{j+1}+\beta'} \in R_j$, it follows that $x_j^{\alpha'}$ is also in R_j. Since $x_j^{p_1 \cdots p_{j+1}\omega_j} = b_j^{-1} \chi_j(F_{j+1,0}) \in R_j$, it follows that h is in the ideal generated by $\chi_j(H_{j+1})$ in R_j. Therefore, $\chi_j(\tilde{K}_j) = L_j \cap R_j$. In particular, $\chi_j(\tilde{K}_j)$ is a prime ideal of R_j and therefore $\tilde{K}_j$ is a prime ideal of A_j. Moreover, there is a chain of graded ring homomorphisms as follows

$$A_j/\tilde{K}_j \cong R_j/(L_j \cap R_j) \hookrightarrow S_j/L_j \cong \mathbb{C}[x_j, x_j^{p_1 \cdots p_j \omega_j}] \hookrightarrow \mathbb{C}[x_j^{\frac{1}{p_{j+1}}}].$$

Let K_{j+1} be the ideal generated by $\tilde{K}_j$ in $A_{j+1} = A_j[y_{j+1}]$ and $x_{j+1} := x_j^{\frac{1}{p_{j+1}}}$. Since $A_{j+1}/K_{j+1} \cong (A_j/\tilde{K}_j)[y_{j+1}]$, it follows that K_{j+1} is a prime ideal of A_{j+1} and there is an injective graded ring homomorphism $\chi_{j+1} : A_{j+1}/K_{j+1} \hookrightarrow \mathbb{C}[x_{j+1}, y_{j+1}] =: S_{j+1}$. Moreover, note that S_j is integral over R_j and $\mathbb{C}[x_{j+1}]$ is integral over $\mathbb{C}[x_j, x_j^{p_1 \cdots p_j \omega_j}]$, it follows that S_{j+1} is integral over $R_{j+1} := \chi_{j+1}(A_{j+1}/K_{j+1})$.

Let $\pi_{j+1} : A_{j+1} \to \mathbb{C}[x, y]$ be as in assertion $8_{(j+1)}$ and J_{j+1} be as in assertion $9_{(j+1)}$, i.e. J_{j+1} is the ideal in A_{j+1} generated by all weighted homogeneous (w.r.t. η_{j+1}) polynomials f such that $\delta_{j+1}(\pi_{j+1}(f)) < \eta_{j+1}(f)$. Since $\delta_{j+1}(\pi_{j+1}(f)) \leq \eta_{j+1}(f)$ for all $f \in A_{j+1}$, it follows that J_{j+1} is a prime ideal of A_{j+1}. Note that $J_j \subseteq J_{j+1}$. Moreover, assertion $4_{(j+1)}$ implies that $H_{j+1} = \mathfrak{L}_{\eta_{j+1}}(F_{j+1})$, so that

$$\begin{aligned}
\delta_{j+1}(\pi_{j+1}(H_{j+1})) &= \delta_{j+1}\left(\pi_{j+1}(F_{j+1}) - \pi_{j+1}(F_{j+1} - H_{j+1})\right) \\
&\leq \max\left\{\delta_{j+1}(f_{j+1}), \delta_{j+1}(\pi_{j+1}(F_{j+1} - H_{j+1}))\right\} \\
&\leq \max\left\{\omega_{j+1}, \eta_{j+1}(F_{j+1} - H_{j+1})\right\} \text{ (due to assertion } 7_{(j+1)}) \\
&< p_{j+1}\omega_j = \eta_{j+1}(H_{j+1}).
\end{aligned}$$

Consequently, $H_{j+1} \in J_{j+1}$, and therefore $K_{j+1} \subseteq J_{j+1}$. We now show that $K_{j+1} = J_{j+1}$. Indeed, assume contrary to the claim that $K_{j+1} \subsetneq J_{j+1}$. Since J_{j+1} is a homogeneous ideal (with respect to the grading induced by η_{j+1}), it follows that $\chi_{j+1}(J_{j+1}/K_{j+1})$ is a non-zero homogeneous prime ideal in R_{j+1} (where R_{j+1} and S_{j+1} are graded by the weighted degree corresponding to weights $\frac{1}{p_1 \cdots p_{j+1}}$ and ω_{j+1} for y_{j+1}). Since S_{j+1} is integral over R_{j+1}, there exists a non-zero homogeneous prime ideal L_{j+1} of S_{j+1} such that $\chi_{j+1}(J_{j+1}/K_{j+1}) = L_{j+1} \cap R_{j+1}$. Since L_{j+1} is homogeneous, either $x_{j+1} \in L_{j+1}$, or $y_{j+1} \in L_{j+1}$, or $y_{j+1}^p - cx_{j+1}^q \in L_{j+1}$ for some non-zero $c \in \mathbb{C}$ and positive integers p, q such that $q/p = p_1 \cdots p_{j+1}\omega_{j+1}$. Consequently, either $x \in J_{j+1}$, or $y_{j+1} \in J_{j+1}$, or $y_{j+1}^{p'} - c^{p'}x^q \in J_{j+1}$, where $p' := p_1 p_2 \cdots p_{j+1}$. It follows straight from the definition of J_{j+1} and η_{j+1} that the first two cases are impossible. Since $\deg_{\mathrm{x}}^{\mathrm{inv}}(f_{j+1}(x, \tilde{\psi}_{j+1}(x, \xi_{j+1}))) = \omega_{j+1}$ (according to assertion $7_{(j+1)}$ of claim 2.19.1), the last case is also impossible. This contradiction shows that $K_{j+1} = J_{j+1}$. In particular, this proves assertion $10_{(j+1)}$ of the claim.

Let $\mathfrak{L}_{\eta_{j+1}}(\ker \pi_{j+1})$ be the ideal in A_{j+1} generated by all weighted homogeneous (w.r.t. η_{j+1}) forms $f \in A_{j+1}$ such that $f = \mathfrak{L}_{\eta_{j+1}}(g)$ for some $g \in \ker \pi_{j+1}$. It follows from the definition that $\mathfrak{L}_{\eta_{j+1}}(\ker \pi_{j+1}) \subseteq J_{j+1}$. On the other hand, for each i, $1 \leq i \leq j+1$, $\tilde{F}_i := F_i - y_i \in \ker \pi_{j+1}$. Since $\eta_j(F_i) = p_i\omega_{i-1} > \omega_i = \eta_j(y_i)$, it follows that $\mathfrak{L}_{\eta_{j+1}}(\tilde{F}_i) = \mathfrak{L}_{\eta_{j+1}}(F_i) = y_{i-1}^{p_i} + F_{i,0} \in \mathfrak{L}_{\eta_{j+1}}(\ker \pi_{j+1})$ for each i, $1 \leq i \leq j+1$. Since $J_{j+1} = K_{j+1}$, it follows by definition of K_{j+1} and inductive assumption $9_{(k)}$ for $k = j$ that J_{j+1} is generated by $y_{i-1}^{p_i} + F_{i,0}$ for $1 \leq i \leq j+1$. Therefore, $J_{j+1} \subseteq \mathfrak{L}_{\eta_{j+1}}(\ker \pi_{j+1})$, and consequently, $\mathfrak{L}_{\eta_{j+1}}(\ker \pi_{j+1}) = J_{j+1}$. It follows that for all $f \in \mathbb{C}[x, y]$,

$$\delta_{j+1}(f) = \min\{\eta_{j+1}(g) : g \in A_{j+1},\ \pi_{j+1}(g) = f\}.$$

In other words, δ_{j+1} is precisely the degree-like function on $\mathbb{C}[x, y]$ induced by η_{j+1}, which proves assertion $8_{(j+1)}$.

It remains to prove assertion $9_{(j+1)}$. For each i, $1 \leq i \leq j+1$, let $H_i := y_{i-1}^{p_i} + F_{i,0}$. We have shown that $\mathcal{B}_{j+1} := (H_{j+1}, \ldots, H_1)$ is a basis of J_{j+1}. To show that $\mathcal{B}_{j+1}$ is a Gröbner basis of J_{j+1} with respect to $\prec_{j+1}$, it suffices to show that running one step of Buchberger's algorithm with $\mathcal{B}_{j+1}$ as input leaves $\mathcal{B}_{j+1}$ unchanged. Here we follow Buchberger's algorithm as described in [**?**, Section 7, Chapter 2]. So pick i, k such that $1 \leq i < k \leq j+1$. Then

$$\mathrm{LT}_{\prec_{j+1}}(H_i) = y_{i-1}^{p_i} \text{ and } \mathrm{LT}_{\prec_{j+1}}(H_k) = y_{k-1}^{p_k}.$$

Therefore, the *S-polynomial* of H_i and H_k is

$$S(H_i, H_k) := y_{k-1}^{p_k} H_i - y_{i-1}^{p_i} H_k = F_{i,0} y_{k-1}^{p_k} - F_{k,0} y_{i-1}^{p_i}.$$

Therefore $S(H_i, H_k)$ is a binomial, and since H_i and H_k are weighted homogeneous forms (with respect to η_{j+1}), it follows that $S(H_i, H_k)$ is also weighted homogeneous and

$$\mathrm{LT}_{\prec_{j+1}}(S(H_i, H_k)) = F_{i,0} y_{k-1}^{p_k}.$$

Next we have to compute the remainder $\overline{S(H_i, H_k)}^{\mathcal{B}_{j+1}}$ of the division of $S(H_i, H_k)$ by $\mathcal{B}_{j+1}$ and adjoin it to $\mathcal{B}_{j+1}$ if it is non-zero.

Assertions $2_{(k)}$, $3_{(k)}$, $2_{(j+1)}$ and $3_{(j+1)}$ imply that for all k', $1 \leq k' \leq j+1$, $\mathrm{LT}_{\prec_{j+1}}(H_{k'}) = y_{k'-1}^{p_{k'}}$ divides $\mathrm{LT}_{\prec_{j+1}}(S(H_i, H_k))$ if and only if $k' = k$. Since

$$\mathrm{LT}_{\prec_{j+1}}(S(H_i, H_k)) = F_{i,0}\,\mathrm{LT}_{\prec_{j+1}}(H_k),$$

the remainder of the division of $S(H_i, H_k)$ by H_k is

$$S_1 := S(H_i, H_k) - F_{i,0}H_k = -F_{k,0}y_{i-1}^{p_i} - F_{i,0}F_{k,0} = -F_{k,0}H_i.$$

Note that for all i' with $i \leq i' \leq j+1$,

$$\deg_{y_{i'}}(S_1) = \deg_{y_{i'}}(F_{k,0}) + \deg_{y_{i'}}(H_i) = \deg_{y_{i'}}(F_{k,0}) < p_{i'+1},$$

and therefore $\mathrm{LT}_{\prec_{j+1}}(H_{i'})$ does *not* divide $\mathrm{LT}_{\prec_{j+1}}(S_1)$. Recall that in the ordering of $\mathcal{B}_{j+1}$, H_p comes *before* H_q if and only if $p > q$. Therefore, the first element of $\mathcal{B}_{j+1}$ whose leading term (w.r.t. $\prec_{j+1}$) divides $\mathrm{LT}_{\prec_{j+1}}(S_1)$ is H_i. Since H_i divides S_1, it follows that the remainder of the division of S_1 by H_i is zero. In particular, $\overline{S(H_i, H_k)}^{\mathcal{B}_{j+1}} = 0$.

Recall that one step of Buchberger's algorithm with input $\mathcal{B}_{j+1}$ consists of adjoining $\overline{S(H_i, H_k)}^{\mathcal{B}_{j+1}}$ to $\mathcal{B}_{j+1}$ for all i, k with $1 \leq i < k \leq j+1$ such that $\overline{S(H_i, H_k)}^{\mathcal{B}_{j+1}}$ is non-zero. Since $\overline{S(H_i, H_k)}^{\mathcal{B}_{j+1}} = 0$ for all i, k with $1 \leq i < k \leq j+1$, running one step of Buchberger's algorithm keeps $\mathcal{B}_{j+1}$ unchanged. It follows that $\mathcal{B}_{j+1}$ is indeed a Gröbner basis of J_{j+1} with respect to $\prec_{j+1}$ (see, e.g. [**?**, Theorem 2, Section 7, Chapter 2]). This completes the proof of assertion $9_{(j+1)}$ and the claim. ☐

3 The Converse

Consider the following statements:

1. There exist degree-wise Puiseux series $\phi_1, \ldots, \phi_m$ in x such that $\phi_j \equiv_r \psi$ for each j, $1 \leq j \leq m$, and $\prod_{j=1}^{m} \Phi_j^* \in \mathbb{C}[x, y]$.
2. ψ satisfies properties $0_{(k)}$ to $10_{(k)}$ for $1 \leq k \leq l$.
3. δ is finitely generated.

We have shown that if δ is positive on non-constant polynomials, then assertion 1 ⇒ assertion 2 ⇒ assertion 3.

Now we show that if δ is positive on non-constant polynomials, then assertion 3 ⇒ assertion 1. At first we introduce some notation. Recall that the*Newton polygon* $\mathrm{NP}(f)$ of $f \in \mathbb{C}[x, y]$ is the convex hull of all $(\alpha, \beta) \in \mathbb{Z}^2$ such that the coefficient of $x^\alpha y^\beta$ in the expansion of f is non-zero. For $f \in \mathbb{C}\langle\langle x \rangle\rangle[y]$, we define *left-filled Newton polyhedron* $\mathrm{LNP}(f)$ of f to be the convex hull of the left-ward pointing horizontal lines starting at $(\alpha, \beta) \in \mathbb{Q}^2$ such that the coefficient of $x^\alpha y^\beta$ in the expansion of f is non-zero.



$f := x^3y^4 + xy^3 + (1 - x^5)y^2 + x^4$

If $f \in \mathbb{C}\langle\langle x\rangle\rangle[y]$ and E is a subset of $\mathbb{R}^2$, then we write $f|_E$ for the sum of the monomial terms in the expansion of f whose exponents lie on E.

Notation 3.1. Let E be an edge of a polyhedron $\mathcal{Q} \subseteq \mathbb{R}^2$. Pick any ray v such that v is normal to E and points *outward* with respect to $\mathcal{Q}$. We define the *normal angle* $\eta(E)$ (resp. *normal slope* $\mu(E)$) of E to be the angle that v makes with the positive x-axis (resp. slope of v). In particular, $\mu(E) = \tan(\eta(E))$.



Lemma 3.2. *Let* $\Phi = \prod_{i=1}^{m} \Phi_i$ *with* $\Phi, \Phi_1, \ldots, \Phi_m \in \mathbb{C}\langle\langle x\rangle\rangle[y]$. *Then*

1. $\mathrm{LNP}(\Phi) = \sum_{i=1}^{m} \mathrm{LNP}(\Phi_i)$.
2. *Let* E *be a non-horizontal edge of* $\mathrm{LNP}(\Phi)$. *Then there are unique faces* E_i *of* $\mathrm{LNP}(\Phi_i)$, $1 \le i \le m$, *such that* $E = E_1 + \cdots + E_m$ *and* $\Phi|_E = \prod_{i=1}^{m} \Phi_i|_E$. *Conversely, for each* i, $1 \le i \le m$, *and every non-horizontal edge* E_i *of* $\mathrm{LNP}(\Phi_i)$, *there exists an edge* E *of* $\mathrm{LNP}(\Phi)$ *such that* $\mu(E) = \mu(E_i)$ *and* $E = E_i + \sum_{j \ne i} E_j$ *for unique faces* E_j *of* $\mathrm{LNP}(\Phi_j)$ *for all* j *such that* $j \ne i$ *and* $1 \le j \le m$.

Proof. Let $\mathcal{P} := \sum_{i=1}^{m} \mathrm{LNP}(\Phi_i)$ and E be a non-horizontal edge of $\mathcal{P}$. Then there are unique faces E_i of $\mathrm{LNP}(\Phi_i)$, $1 \le i \le m$, such that $E = E_1 + \cdots + E_m$ and $\Phi|_E = \prod_{i=1}^{m} \Phi_i|_E$. It follows that $\mathcal{P} \subseteq \mathrm{LNP}(\Phi)$. It is straightforward to see that the opposite inclusion is also true, and therefore $\mathrm{LNP}(\Phi) = \mathcal{P}$. The last assertion of the lemma follows similarly. ☐

Example 3.3. Let $f := x^3y^4 + xy^3 + (1 - x^5)y^2 + x^4$. Then f has a factorization in $\mathbb{C}\langle\langle x\rangle\rangle[y]$ of the form

$$f = x^3(y + (x + \text{l.d.t.}))(y + (-x + \text{l.d.t.}))(y + (x^{-\frac{1}{2}} + \text{l.d.t.}))(y + (-x^{-\frac{1}{2}} + \text{l.d.t.})).$$



P_1 := LNP(x^3(y + (x + l.d.t))



P_2 := LNP(y + (-x + l.d.t))





P_3 := LNP(y + ($x^{-1/2}$ + l.d.t))



P_4 := LNP(y + ($-x^{-1/2}$ + l.d.t))

NP(f) = $P_1 + P_2 + P_3 + P_4$

We now assume that δ is positive on non-constant polynomials and go back to the proof of assertion 1 from assertion 3. So assume δ is finitely generated. Let $g_0 := 1$ and pick

$g_1, \dots, g_N \in \mathbb{C}[x,y]$ such that X^δ is isomorphic to the closure in $\mathbb{P}^N$ of the image of $\mathbb{C}^2$ under the mapping

$$\sigma : (x,y) \mapsto (g_0(x,y) : g_1(x,y) : \cdots : g_N(x,y)).$$

W.l.o.g. assume that $\{x,y\} \subseteq \{g_1, \dots, g_N\}$. Let $\tilde{y} := y - \psi(x)$. For each i, $0 \le i \le N$, let $\mathcal{P}_i$ (resp. $\tilde{\mathcal{P}}_i$) be the Newton polygon of g_i with respect to (x,y)-coordinates (resp. left-filled Newton polyhedron of g_i with respect to $(x,\tilde{y})$-coordinates) and let $\mathcal{P}$ (resp. $\tilde{\mathcal{P}}$) be the convex hull of $\bigcup_{0=1}^N \mathcal{P}_i$ (resp. $\bigcup_{0=1}^N \tilde{\mathcal{P}}_i$).

Claim 3.3.1.

1. *Let E be a non-horizontal edge of $\tilde{\mathcal{P}}$ such that $\mu(E) \neq r$. Then $E \subseteq \tilde{\mathcal{P}}_i$ for all i, $1 \le i \le N$, such that $E \cap \tilde{\mathcal{P}}_i \neq \emptyset$.*
2. *There is no edge E of $\mathcal{P}$ such that $\pi > \eta(E) \ge \pi/2$.*

Proof. Note that every vertex v of $\mathcal{P}$ (resp. $\tilde{\mathcal{P}}$) is a vertex of one of the $\mathcal{P}_i$'s (resp. $\tilde{\mathcal{P}}_i$'s). At first we prove assertion 1 by contradiction. So assume that there is a non-horizontal edge E of $\tilde{\mathcal{P}}$ such that $s := \mu(E) \neq r$ and there exists i, $1 \le i \le N$, such that $\emptyset \neq E \cap \tilde{\mathcal{P}}_i \subsetneq E$. W.l.o.g. we may assume $i = 1$. There exists a vertex v of E such that $v \notin \mathcal{P}_1$. W.l.o.g. we may assume that v is a vertex of $\tilde{\mathcal{P}}_2$, $1 \le i \le 2$.



CHANGE PICTURE

For each $c \in \mathbb{C}$, let γ_c be the curve in $\mathbb{C}^2$ given by $y = \psi(x) + cx^s$. Let $M := \alpha + \beta s$ for any $(\alpha, \beta) \in E$. Then for all i, $1 \le i \le N$, and all $t \in \mathbb{C} \setminus \{0\}$,

$$g_i(\gamma_c(t)) = g_i|_E(1,c)t^M + \text{l.d.t.}\ ,$$

where we wrote $g_i|_E$ for the sum of the monomial terms in the expansion of g_i (in $(x,\tilde{y})$-coordinates) whose exponents lie on E. Recall that $(1,0) \in \tilde{\mathcal{P}}$ by our assumption on g_i's. In particular, this implies that $M \ge 1$. It follows that

$$\begin{aligned}\lim_{t\to\infty} \sigma(\gamma_c(t)) &= \lim_{t\to\infty} (1 : g_1(\gamma_c(t)) : g_2(\gamma_c(t)) : \cdots : g_N(\gamma_c(t))) \\ &= \lim_{t\to\infty} (0 : g_1|_E(1,c) : g_2|_E(1,c) : \cdots : g_N|_E(1,c))\end{aligned}$$

Since E is non-horizontal, our choice of $\tilde{\mathcal{P}}_1$ and $\tilde{\mathcal{P}}_2$ implies that $\frac{g_2|_E}{g_1|_E}(1,c)$ is a non-constant rational function of c. Therefore, for generic $c \in \mathbb{C}$, $\lim_{t\to\infty} \sigma(\gamma_c(t))$ is a generic point of

$X_\infty := X^\delta \setminus \mathbb{C}^2$. But then lemma **??** implies that for almost all $c \in \mathbb{C}$, the degree-wise Puiseux series for γ_c is of the form:

$$y = \zeta^{k_c} \star_{p_1 \cdots p_l} \psi(x) + h(c)x^r + \text{l.d.t.}$$

for a non-constant function h of c and some $k_c \in \mathbb{Z}$ (recall that ζ is a $p_1 \cdots p_l$-th root of unity). In particular, the coefficient of x^r in the degree-wise Puiseux series for γ_c is non-zero. This gives a contradiction to our choice of γ_c and proves assertion 1.

Now we prove assertion 2 by contradiction. So assume $\mathcal{P}$ has an edge E such that $\eta(E) \in [\pi/2, \pi)$. Let $s := 1/\mu(E)$. Then $s \geq 0$ and $(s, 1)$ is an outward pointing normal to E. Let the vertices of E be v_1 and v_2. Renumbering g_i's if necessary, we may assume that there exists $k \leq N$ such that $\mathcal{P}_i \cap E \neq \emptyset$ if and only if $1 \leq i \leq k$. Moreover, w.l.o.g. we may also assume that either

Case 1. E is a vertex of $\mathcal{P}_i$ for all i, $1 \leq i \leq k$, or

Case 2. v_i is a vertex of $\mathcal{P}_i$, $1 \leq i \leq 2$.



Case 1



Case 2

For each $c \in \mathbb{C}$, let γ_c be the curve in $\mathbb{C}^2$ given by $\gamma_c(t) := (ct^s, t)$. Let $M := \alpha s + \beta$ for any $(\alpha, \beta) \in E$. Then for all i, $1 \leq i \leq N$, and all $t \in \mathbb{C} \setminus \{0\}$,

$$g_i(\gamma_c(t)) = g_i|_E(c, 1)t^M + \text{l.d.t.}$$

Since y appears as one of the g_i's, it follows that $(0, 1) \in \mathcal{P}$. In particular, this implies that $M \geq 1$. It follows that

$$\begin{aligned}\lim_{t\to\infty} \sigma(\gamma_c(t)) &= \lim_{t\to\infty} (1 : g_1(\gamma_c(t)) : g_2(\gamma_c(t)) : \cdots : g_N(\gamma_c(t))) \\ &= \lim_{t\to\infty} (0 : g_1|_E(c, 1) : g_2|_E(c, 1) : \cdots : g_k|_E(c, 1) : 0 : \cdots : 0).\end{aligned}$$

Note that E is *not* vertical. Therefore different integral points on E have different x-coordinates. Consequently, if there exists i, j such that $1 \leq i \leq j \leq k$ and $\frac{g_i|_E}{g_j|_E}$ is *not* constant, then $\frac{g_i|_E}{g_j|_E}(c, 1)$ is a non-constant rational function of c, and therefore for generic $c \in \mathbb{C}$, $\lim_{t\to\infty} \sigma(\gamma_c(t))$ is a generic point of X_∞. As in the proof of assertion 1, an application of lemma **??** then yields a contradiction. In particular, this implies that Case 2 does not occur.

So assume that for all i, $1 \leq i \leq k$, $g_i|_E = \lambda_i g_1|_E$ for some $\lambda_i \in \mathbb{C}$. Then the situation is that of Case 1. In particular, $g_1|_E$ is *not* a monomial. It follows that there is a factor of g_1 of the form $x^p y^q - b$ for some $b \neq 0 \in \mathbb{C}$ and integers p, q such that either $(p, q) = (1, 0)$ (in the case that E is horizontal), or p, q are relatively prime positive integers such that $\frac{q}{p} = -s$. In particular, this implies that there exists a degree-wise Puiseux series $\phi(y) = \sum_{j \geq 1} b_j y^{s_j} \in \mathbb{C}\langle\langle y \rangle\rangle$ such that $s = s_1 > s_2 > \cdots$, $b_1 := b^{1/p}$ and $x - \phi(y)$ divides g_1 in $\mathbb{C}\langle\langle y \rangle\rangle[x]$. We now recall, following [**?**, Chapter IV, Section 3], the algorithm for computing ϕ:

Assume we have computed the first j terms of ϕ. Let x_j be an indeterminate and set

$$g_{1,j}(x_j, y) := g_1(\sum_{i=1}^{j} b_i y^{s_i} + x_j, y).$$

If x_j divides $g_{1,j}$, then set $\phi := \sum_{i=1}^{j} b_i y^{s_i}$ and stop here. Otherwise, there exists a rational number ω such that

$\omega < s_j$ and the leading form $G_{1,j,\omega}$ of $g_{1,j}$ with respect to the weighted degree corresponding to weights $(\omega, 1)$ for (x_j, y) is *not* a monomial. $(P_{\omega,j})$

Let s_{j+1} be the *largest* rational number such that $\omega = s_{j+1}$ satisfiese $(P_{\omega,j})$ and b_{j+1} to be a non-zero complex number such that $x_j - b_{j+1} y^{s_{j+1}}$ divides $G_{1,j,s_{j+1}}$.

Let $x_0 := x$ and n be the polydromy order of ϕ. For each $j \geq 0$, let ω_j be the weighted degree on $\mathbb{C}(x_j, y^{1/n})$ corresponding to weights $(s_{j+1}, 1)$ for (x_j, y), and define

$$\begin{gathered}
\phi_j := \sum_{i=1}^{j} b_i y^{s_i}, \\
g_{i,j}(x_j, y) := g_i(\phi_j(y) + x_j, y),\ d_{i,j} := \omega_j(g_{i,j}),\ G_{i,j}(x_j, y) := \mathfrak{L}_{\omega_j}(g_{i,j}),\ 1 \leq i \leq N, \\
M_j := \max\{d_{i,j} : 1 \leq i \leq N\}, \text{ and} \\
\mathcal{I}_j := \{i : 1 \leq i \leq N \text{ and } d_{i,j} = M_j\}.
\end{gathered}$$

Note that each $G_{i,j} \in \mathbb{C}[x_j, y^{1/n}, y^{-1/n}]$. Moreover, for $j = 0$ we see that $M_0 = M$, $\mathcal{I}_0 = \{1, \ldots, k\}$, and $G_{i,0} = g_i|_E$ for each $i \in \mathcal{I}_0$. Recall that there exists i^*, $1 \leq i^* \leq N$, such that $g_{i^*} = y$. In particular, $M_j \geq d_{i^*,j} = 1$ for all $j \geq 0$. Since $\lim_{j \to \infty} d_{1,j} = -\infty$ (because $x = \phi(y)$ is a root of g_1), it follows that there exists $j^* \geq 1$ such that $1 \in \mathcal{I}_{j^*-1}$ and $1 \notin \mathcal{I}_j$ for all $j \geq j^*$.

For each i, $1 \leq i \leq N$, let $\mathcal{P}_i^*$ be the Newton polygon of g_{i,j^*}, and let $\mathcal{P}^*$ be the convex hull of $\bigcup_{i=1}^{N} \mathcal{P}_i^*$. Let $\mathcal{E}^*$ be the collection of all edges E^* of $\mathcal{P}^*$ such that there exists $s^* \in [s_{j^*+1}, s_{j^*}]$ such that $(s^*, 1)$ is an *outward pointing* normal to E^*.

Claim 3.3.2. *There exists $E^* \in \mathcal{E}^*$ and i^*, $1 < i^* \leq N$ such that $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}$ is a non-constant rational function in (x_{j^*}, y).*

Proof. For each $s^* \in \mathbb{Q}$, let $\omega^*_{s^*}$ be the weighted degree on $\mathbb{C}(x_{j^*}, y^{1/n})$ corresponding to weights $(s^*, 1)$ for (x_{j^*}, y). Abusing the notation, we also write $\omega^*_{s^*}$ for the linear functional on $\mathbb{R}^2$ which maps $(\alpha, \beta) \mapsto \alpha s^* + \beta$. For each $s^* \in \mathbb{Q}$, let $F^*_{s^*}$ be the maximal face (i.e. either an edge, or a vertex) of $\mathcal{P}^*$ where $\omega^*_{s^*}|_{\mathcal{P}^*}$ achieves the maximum. Note that for all i, $1 \leq i \leq N$,

$$\omega^*_{s_{j^*}}(g_{i,j^*}) = \omega_{j^*-1}(g_{i,j^*-1}), \text{ and} \tag{25}$$

$$\omega^*_{s_{j^*+1}}(g_{i,j^*}) = \omega_{j^*}(g_{i,j^*}) \tag{26}$$

Let $\mathcal{V}$ be the collection of vertices v of $\mathcal{P}^*$ such that $v = F^*_{s^*}$ for some $s^* \in [s_{j^*+1}, s_{j^*}]$ and $\mathcal{V}^* := \{v \in \mathcal{V} : v \not\in \mathcal{P}^*_1\}$. Identity (26) implies that $\mathcal{V}^* \neq \emptyset$. Order the elements in $\mathcal{V}$ *counterclockwise*, i.e. $v_1 \prec v_2$ iff $v_1 \neq v_2$ and the ordering O, v_1, v_2 (where O is the origin of $\mathbb{R}^2$) gives a counterclockwise orientation of the triangle determined by O, v_1 and v_2. Let v^* be the first (i.e. the smallest) element of $\mathcal{V}^*$ with respect to $\prec$. Let $\mathcal{V}^{**} := \{v \in \mathcal{V} \setminus \mathcal{V}^* : v \prec v^*\}$. There are two possible scenarios:

Scenario 1. $\mathcal{V}^{**} \neq \emptyset$. In this case let v be the maximal element of $\mathcal{V}^*$ and E^* be the edge of $\mathcal{P}^*$ with vertices v and v^*.

Scenario 2. v^* is the smallest element of $\mathcal{V}$. In this case identity (25) implies that $F^*_{s_{j^*}}$ is an edge of $\mathcal{P}^*$ and v^* is a vertex of $F^*_{s_{j^*}}$. Set $E^* := F^*_{s_{j^*}}$.

It follows from the definition of E^* and v^* that $E^* \in \mathcal{E}^*$, $E^* \cap \mathcal{P}^*_1 \neq \emptyset$, v^* is a vertex of E^*, and $v^* \not\in \mathcal{P}^*_1$. Pick i^*, $2 \leq i \leq N$, such that $v^* \in \mathcal{P}^*_{i^*}$. Then $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}$ is a non-constant rational function in (x_{j^*}, y), as required. ☐

Let E^* be as in claim 3.3.2 and $s^* \in [s_{j^*+1}, s_{j^*}]$ such that $E^* = F^*_{s^*}$. For each $c \in \mathbb{C}$, let γ^*_c be the curve in $\mathbb{C}^2$ given by

$$\gamma^*_c(t) := (\phi_{j^*}(t) + ct^{s^*}, t).$$

Let $M^* := \omega^*_{s^*}(x^\alpha_{j^*} y^\beta)$ for any $(\alpha, \beta) \in E^*$. Then for all i, $1 \leq i \leq N$, and all $t \in \mathbb{C} \setminus \{0\}$,

$$g_i(\gamma^*_c(t)) = g_{i,j^*}|_{E^*}(c, 1)t^{M^*} + \text{l.d.t.}$$

Let i^* be as in claim 3.3.2. Then $g_{1,j^*}|_{E^*}$ and $g_{i^*,j^*}|_{E^*}$ are weighted homogeneous with respect to $\omega^*_{s^*}$ with weight M^* and claim 3.3.2 implies that $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}(c, 1)$ is a non-constant rational function of c. It then follows in the same way as in the proof of assertion 1 that for generic $c \in \mathbb{C}$, $\lim_{t\to\infty} \sigma(\gamma^*_c(t))$ is a generic point of X_∞, and therefore lemma **??** implies that for almost all $c \in \mathbb{C}$, the degree-wise Puiseux series $\phi^*_c(y)$ for x (in terms of y) associated to γ^*_c agrees up to the *generic term* with the degree-wise Puiseux series $\psi^*(y)$ for x associated to δ. But $\deg_y(\psi^*(y)) = \delta(x)/\delta(y) > 0$, whereas $\deg_y(\phi^*_c(y)) = \deg_y(\phi_{j^*}(t) + ct^{s^*}) = s_1 \leq 0$, which is a contradiction. This contradiction proves assertion 2 and finishes the proof of claim 3.3.1. ☐

We now go back to the proof of assertion 1 from assertion 3. For each i, $1 \leq i \leq N$, let

$$g_i = x^{\alpha_i} y^{\beta_i} \prod_{j=1}^{m_i} \Phi^*_{i,j} \tag{27}$$

be the factorization of g_i in $\mathbb{C}\langle\langle x\rangle\rangle[y]$, where for each j, $1 \leq j \leq m_i$,

$$\Phi^*_{i,j} := \prod_{k=0}^{m_{i,j}-1} (y - \zeta_{i,j}^k \star_{m_{i,j}} \phi_{i,j}(x)) \tag{28}$$

where $\phi_{i,j}(x) \in \mathbb{C}\langle\langle x\rangle\rangle$ with polydromy order $m_{i,j}$ and $\zeta_{i,j}$ is an $m_{i,j}$-th root of unity. Assume to the contrary of the claim that assertion 1 is not true. Then for each i, $1 \leq i \leq N$, either

i-1. $\alpha_i > 0$, or,

i-2. renumbering $\phi_{i,j}$'s if necessary, we may assume that for each k, $0 \leq k < m_{i,0}$,

$$s_{i,k} := \deg_x \Big(\psi(x) - \zeta_{i,0}^k \star_{m_{i,0}} \phi_{i,0}(x)\Big) > r.$$

In this case, replacing $\phi_{i,0}$ by $\zeta_{i,0}^k \star_{m_{i,0}} \phi_{i,0}(x)$ for some k, $0 \leq k < m_{i,0}$, we may (and will) also assume that $s_{i,0} = \min\{s_{i,k} : 0 \leq k < m_{i,0}\}$.

Let $\tilde{y}$, $\tilde{\mathcal{P}}$ and $\tilde{\mathcal{P}}_1, \ldots, \tilde{\mathcal{P}}_N$ be as in claim 3.3.1. We claim that $\mathcal{P}$ has a non-horizontal edge E such that $\mu(E) > r$. Indeed, pick any i, $1 \leq i \leq N$, such that $g_i \notin \mathbb{C}[x]$. Then either $\beta_i > 0$ or $m_i > 0$, where β_i and m_i are as in (27). Note that $y = \tilde{y} + \psi(x) = y + a_1 x^{q_1/p_1} + \text{l.d.t.}$. Therefore the left-filled Newton polygon of y in $(x, \tilde{y})$-coordinates has only one non-horizontal edge E_y with $\mu(E_y) = q_1/p_1$.



LNP($\tilde{y} + \psi(x)$)

Similarly, for each j, $1 \leq j \leq m_i$, there is at least one k, $0 \leq k \leq m_{i,j} - 1$, such that the only non-horizontal edge E of the left-filled Newton polygon (in $(x, \tilde{y})$-coordinates) of $y - \zeta_{i,j}^k \star_{m_{i,j}} \phi_{i,j}(x) = \tilde{y} + (\psi(x) - \zeta_{i,j}^k \star_{m_{i,j}} \phi_{i,j}(x))$ satisfies $\mu(E) \geq q_1/p_1$. It follows then from assertion 2 of lemma 3.2 that there is an edge E_i of $\tilde{\mathcal{P}}_i$ such that $\mu(E_i) \geq q_1/p_1 > r$. In particular, if E_i is the *top-most* edge of $\tilde{\mathcal{P}}_i$, then $\mu(E_i) > r$. Now, let $\tilde{v}$ be the top-most vertex of $\tilde{\mathcal{P}}$. Then there exists i, $1 \leq i \leq N$, such that $\tilde{v}$ is the top-most vertex of $\tilde{\mathcal{P}}_i$ as well. Then assertion 1 of claim 3.3.1 implies that the top-most edge $\tilde{E}$ of $\tilde{\mathcal{P}}$ is also the top-most edge of $\tilde{\mathcal{P}}_i$. It then follows from the preceding discussion that $\mu(\tilde{E}) > r$, as required.

Let $\tilde{v}$ be as in the preceding paragraph. Renumbering g_i's if necessary, we may assume that $\tilde{v} \in \tilde{\mathcal{P}}_1$. We claim that $\alpha_1 = 0$, where α_1 is as in (27). Indeed, since $\tilde{v}$ is the top-most point of $\tilde{\mathcal{P}}_1$, it follows that $\tilde{v} = (\alpha_1, \beta_1 + \sum_{j=1}^{m_1} m_{1,j})$. Now, note that the change of coordinates $(x, y) \mapsto (x, \tilde{y})$ does *not* change the coordinates of the top-most vertex of the Newton polygon (or left-filled Newton polyhedron) of any rational function. It follows that the top-most point of $\mathcal{P}$ also has the same coordinates as that of $\tilde{v}$. In particular, the top-most point of $\mathcal{P}$ is *not* on y-axis. Since $(0, 1) \in \mathcal{P}$ (because there exists i, $1 \leq i \leq N$, such that $g_i = y$), it follows that there is an edge E of $\mathcal{P}$ such that $\eta(E) \in [\pi/2, \pi)$. This contradicts assertion 2 of claim 3.3.1 and shows that $\alpha_1 = 0$.

The preceding paragraph implies that $i = 1$ satisfies property i-2. In particular, $\psi(x) - \phi_{1,0}(x) = b_{1,0}x^{s_{1,0}} +$ l.d.t. for some $b_{1,0} \neq 0 \in \mathbb{C}$ and the non-horizontal edge E_0 of LNP$(\tilde{y} + \psi(x) - \phi_{1,0}(x))$ satiesfies: $\mu(E_0) = s_{1,0}$, and

$$(\tilde{y} + \psi(x) - \phi_{1,0}(x))|_{E_0} = \tilde{y} + b_{1,0}x^{s_{1,0}}.$$

Lemma 3.2 implies that $\tilde{\mathcal{P}}_1$ has an edge $\tilde{E}$ such that $\mu(\tilde{E}_1) = s_{1,0}$ and

$$g_1|_{\tilde{E}_1} = (\tilde{y} + b_{1,0}x^{s_{1,0}})h_1 \tag{29}$$

for some $h_1 \in \mathbb{C}[\tilde{y}, x^{\frac{1}{p_{1,0}}}, x^{-\frac{1}{p_{1,0}}}]$ where $s_{1,0} = \frac{q_{1,0}}{p_{1,0}}$ with $p_{1,0}, q_{1,0} \in \mathbb{Z}$. It follows that there is $\phi(x) = \sum_{j \geq 1} b_j x^{s_j} \in \mathbb{C}\langle\langle x \rangle\rangle$ such that

$$\begin{gathered} s_{1,0} = s_1 > s_2 > \cdots, \\ b_1 := -b_{1,0}, \text{ and} \\ \tilde{y} - \phi(x) \text{ divides } g_1 \text{ in } \mathbb{C}\langle\langle x \rangle\rangle[\tilde{y}]. \end{gathered}$$

Let n be the polydromy order of ϕ. For each non-negative integer j, let $\tilde{y}_j$ be an indeterminate and ω_j be the weighted degree on $\mathbb{C}(x^{1/n}, \tilde{y}_j)$ corresponding to weights $(1, s_{j+1})$ for $(x, \tilde{y}_j)$, and define

$$\begin{gathered} \phi_j := \sum_{i=1}^{j} b_i x^{s_i}, \\ g_{i,j}(x, \tilde{y}_j) := g_i(x, \tilde{y}_j + \phi_j(x)),\ d_{i,j} := \omega_j(g_{i,j}),\ G_{i,j}(x, \tilde{y}_j) := \mathfrak{L}_{\omega_j}(g_{i,j}),\ 1 \leq i \leq N, \\ M_j := \max\{d_{i,j} : 1 \leq i \leq N\}, \text{ and} \\ \mathcal{I}_j := \{i : 1 \leq i \leq N \text{ and } d_{i,j} = M_j\}. \end{gathered}$$

Note that each $G_{i,j} \in \mathbb{C}[\tilde{y}_j, x^{1/n}, x^{-1/n}]$. Also, recall that there exists i, $1 \leq i \leq N$, such that $g_i = x$. It follows that $M_j \geq \omega_j(g_{i,j}) = \omega_j(x) = 1$ for all $j \geq 0$.

Now, recall that the top-most edges of $\tilde{\mathcal{P}}$ and $\tilde{\mathcal{P}}_1$ coincide and has normal slope greater than r. It then follows due to assertion 1 of claim 3.3.1 that the set of non-horizontal edges E of $\tilde{\mathcal{P}}$ such that $\mu(E) > r$ is identical to the set of non-horizontal edges E' of $\tilde{\mathcal{P}}_1$ such that $\mu(E') > r$. In particular, $\tilde{E}_1$ is also an edge of $\tilde{\mathcal{P}}$, which implies that $1 \in \mathcal{I}_0$.

PICTURREEE

Since $y = \phi(x)$ is a root of g_1, it follows that $\lim_{j \to \infty} d_{1,j} = -\infty$. Since $M_j \geq 1$ for all $j \geq 0$, it follows that $d_{1,j} < M_j$ for sufficiently large j. Therefore there exists $j^* \geq 1$ such that $1 \in \mathcal{I}_{j^*-1}$ and $1 \notin \mathcal{I}_{j^*}$.

For each i, $1 \leq i \leq N$, let $\tilde{\mathcal{P}}^*_i$ be the left-filled Newton polyhedron of g_{i,j^*}, and let $\tilde{\mathcal{P}}^*$ be the convex hull of $\bigcup_{i=1}^N \tilde{\mathcal{P}}^*_i$. Let $\mathcal{E}^*$ be the collection of all edges E^* of $\tilde{\mathcal{P}}^*$ such that there exists $s^* \in [s_{j^*+1}, s_{j^*}]$ such that $(1, s^*)$ is an *outward pointing* normal to E^*.

Claim 3.3.3. *There exists $E^* \in \mathcal{E}^*$ and i^*, $1 < i^* \leq N$ such that $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}$ is a non-constant rational function in $(x, \tilde{y}_{j^*})$.*

Proof. For each $s^* \in \mathbb{Q}$, let $\omega^*_{s^*}$ be the weighted degree on $\mathbb{C}(x^{1/n}, \tilde{y}_{j^*})$ corresponding to weights $(1, s^*)$ for $(x, \tilde{y}_{j^*})$. Abusing the notation, we also write $\omega^*_{s^*}$ for the linear functional on $\mathbb{R}^2$ which maps $(\alpha, \beta) \mapsto \alpha + \beta s^*$. For each $s^* \in \mathbb{Q}$, let $F^*_{s^*}$ be the maximal face (i.e. either an edge, or a vertex) of $\tilde{\mathcal{P}}^*$ where $\omega^*_{s^*}|_{\tilde{\mathcal{P}}^*}$ achieves the maximum. Note that for all i, $1 \leq i \leq N$,

$$\omega^*_{s_{j^*}}(g_{i,j^*}) = \omega_{j^*-1}(g_{i,j^*-1}), \text{ and} \tag{30}$$

$$\omega^*_{s_{j^*+1}}(g_{i,j^*}) = \omega_{j^*}(g_{i,j^*}) \tag{31}$$

Let $\mathcal{V}$ be the collection of vertices v of $\tilde{\mathcal{P}}^*$ such that $v = F^*_{s^*}$ for some $s^* \in [s_{j^*+1}, s_{j^*}]$ and $\mathcal{V}^* := \{v \in \mathcal{V} : v \notin \tilde{\mathcal{P}}^*_1\}$. Since $1 \notin \mathcal{I}_{j^*}$, identity (26) implies that $\mathcal{V}^* \neq \emptyset$. Order the elements in $\mathcal{V}$ *counterclockwise*, i.e. $v_1 \prec v_2$ iff $v_1 \neq v_2$ and the ordering O, v_1, v_2 (where O is the origin of $\mathbb{R}^2$) gives a counterclockwise orientation of the triangle determined by O, v_1 and v_2. Let v^* be the smallest element of $\mathcal{V}^*$ with respect to $\prec$ and define $\mathcal{V}^{**} := \{v \in \mathcal{V} \setminus \mathcal{V}^* : v \prec v^*\}$. There are two possible scenarios:

Scenario 1. $\mathcal{V}^{**} \neq \emptyset$. In this case let v be the maximal element of $\mathcal{V}^*$ and E^* be the edge of $\tilde{\mathcal{P}}^*$ with vertices v and v^*.

Scenario 2. $\mathcal{V}^{**} = \emptyset$. In this case v^* is the smallest element of $\mathcal{V}$. Since $1 \in \mathcal{I}_{j^*-1}$, identity (25) then implies that $F^*_{s_{j^*}}$ is an edge of $\tilde{\mathcal{P}}^*$ and v^* is a vertex of $F^*_{s_{j^*}}$. Set $E^* := F^*_{s_{j^*}}$.

It follows from the definition of E^* and v^* that $E^* \in \mathcal{E}^*$, $E^* \cap \tilde{\mathcal{P}}^*_1 \neq \emptyset$, v^* is a vertex of E^*, and $v^* \notin \tilde{\mathcal{P}}^*_1$. Pick i^*, $2 \leq i \leq N$, such that $v^* \in \tilde{\mathcal{P}}^*_{i^*}$. Then $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}$ is a non-constant rational function in $(x, \tilde{y}_{j^*})$, as required. □

Let E^* be as in claim 3.3.3 and $s^* \in [s_{j^*+1}, s_{j^*}]$ such that $E^* = F^*_{s^*}$. For each $c \in \mathbb{C}$, let γ^*_c be the curve in $\mathbb{C}^2$ given by

$$(x(t), \tilde{y}(t)) := (t, \phi_{j^*}(t) + ct^{s^*}).$$

Let $M^* := \omega^*_{s^*}(x^\alpha \tilde{y}^\beta_{j^*})$ for any $(\alpha, \beta) \in E^*$. Then for all i, $1 \leq i \leq N$, and all $t \in \mathbb{C} \setminus \{0\}$,

$$g_i(\gamma^*_c(t)) = g_{i,j^*}|_{E^*}(1, c)t^{M^*} + \text{l.d.t.}$$

Let i^* be as in claim 3.3.3. Then $g_{1,j^*}|_{E^*}$ and $g_{i^*,j^*}|_{E^*}$ are weighted homogeneous with respect to $\omega^*_{s^*}$ with weight M^* and claim 3.3.3 implies that $\frac{g_{i^*,j^*}|_{E^*}}{g_{1,j^*}|_{E^*}}(1, c)$ is a non-constant rational function of c. It then follows in the same way as in the proof of assertion 1 of claim 3.3.1 that for generic $c \in \mathbb{C}$, $\lim_{t \to \infty} \sigma(\gamma^*_c(t))$ is a generic point of X_∞, and therefore lemma **??** implies that for almost all $c \in \mathbb{C}$, the degree-wise Puiseux series $\phi^*_c(x)$ for y (in terms of x) is of the form:

$$\phi^*_c(x) = \zeta^{k_c} \star_{p_1 \cdots p_l} \psi(x) + h(c)x^r + \text{l.d.t.}$$

for a non-constant function h of c and $k_c \in \mathbb{Z}$, $0 \leq k_c < p_1 \cdots p_l$. On the other hand, it follows from the definition of γ^*_c that

$$\phi^*_c(x) = \tilde{y}(x) + \psi(x) = \psi(x) + \phi_{j^*}(x) + cx^{s^*} = \psi(x) + (-b_{1,0}x^{s_{1,0}} + \text{l.d.t.}).$$

It follows that

$$\begin{aligned}
&\deg_x \Big(\psi(x) - b_{1,0}x^{s_{1,0}} - \zeta^{k_c} \star_{p_1\cdots p_l} \psi(x)\Big) < s_{1,0}\\
\Rightarrow &\tilde{s}_c := \deg_x \Big(\phi_{1,0}(x) - \zeta^{k_c} \star_{p_1\cdots p_l} \psi(x)\Big) < s_{1,0}.
\end{aligned}$$

Let the truncation of ψ at $\tilde{s}_c$ be

$$\psi_{\tilde{s}_c} = \sum_{i=1}^{n-1}\sum_{j=0}^{k_i} a_{i,j}x^{\frac{q_{i,j}}{p_1\cdots p_i}} + \sum_{j=0}^{k'_n} a_{n,j}x^{\frac{q_{n,j}}{p_1\cdots p_n}},\ \text{so that}$$

$$\frac{q_{n,j}}{p_1\cdots p_n} > \tilde{s}_c \geq \deg_x \Big(\phi_{1,0}(x) - \zeta^{k_c} \star_{p_1\cdots p_l} \psi_{\tilde{s}_c}(x)\Big) = \deg_x \Big(\phi_{1,0}(x) - \zeta_n^{k_c} \star_{p_1\cdots p_n} \psi_{\tilde{s}_c}(x)\Big)$$

It follows that the first n Puiseux pairs of $\phi_{1,0}$ are (p_i, q_i), $1 \leq i \leq n$. In particular, $p_1 \cdots p_n$ divides $m_{1,0}$ (which is the the polydromy order of $\phi_{1,0}$) and therefore $a \star_{m_{1,0}} \psi_{\tilde{s}_c}$ is well defined for all $a \in \mathbb{C}$. Since

$$\begin{aligned}
\zeta_{1,0}^{-k_c} \star_{m_{1,0}} \Big(\phi_{1,0}(x) - \zeta_n^{k_c} \star_{p_1\cdots p_n} \psi_{\tilde{s}_c}(x)\Big) &= \zeta_{1,0}^{-k_c} \star_{m_{1,0}} \phi_{1,0}(x) - \zeta_{1,0}^{-k_c} \star_{m_{1,0}} \Big(\zeta_n^{k_c} \star_{p_1\cdots p_n} \psi_{\tilde{s}_c}(x)\Big)\\
&= \zeta_{1,0}^{-k_c} \star_{m_{1,0}} \phi_{1,0}(x) - \zeta_n^{-k_c} \star_{p_1\cdots p_n} \Big(\zeta_n^{k_c} \star_{p_1\cdots p_n} \psi_{\tilde{s}_c}(x)\Big)\\
&= \zeta_{1,0}^{m_{1,0}-k_c} \star_{m_{1,0}} \phi_{1,0}(x) - \psi_{\tilde{s}_c}(x),
\end{aligned}$$

it follows that

$$s_{1,m_{1,0}-k_c} = \deg_x \Big(\psi(x) - \zeta_{1,0}^{m_{1,0}-k_c} \star_{m_{1,0}} \phi_{1,0}(x)\Big) \leq \tilde{s}_c < s_{1,0}.$$

This contradicts the minimality of $s_{1,0}$ and proves that assertion 1 is indeed true.

4 Old Stuff

Let $\bar{X}$ be a normal algebraic surface containing $S := \mathbb{C}^2$ as a dense open subset and V be an irreducible component of $X_\infty := \bar{X} \setminus X$. Then $\dim V = 1$ (see, e.g., [2, Proposition 1]). Let δ be the semidegree associated to V. Then δ is divisorial. Moreover, since δ corresponds to a semidegree 'at infinity', there is $f \in \mathbb{C}[x, y]$ such that $\delta(f) > 0$. Therefore, making a linear change of coordinates if necessary, we may assume that $\delta(x) > 0$. Then according to corollary 2.7, there is a finite Puiseux series ψ in x and a rational number $r < \mathrm{ord}_x(\psi)$ such that (2) holds for all $f \in \mathbb{C}[x, y]$.

Let p be the polydromy order of ψ and for all j, $0 \leq j \leq p-1$, let ψ_j be the j-th conjugate of ψ. Define

$$\Psi := \prod_{j=0}^{p-1}(y - \psi_j).$$

Then $\Psi \in \mathbb{C}[x, x^{-1}, y]$. Let Ψ_1 be the sum of the monomials in Ψ with δ-value greater than $\delta(\Psi)$ and $\Psi_2 := \Psi - \Psi_1$. Note that

1. Ψ_1 is always non-zero, since y^p is a monomial in Ψ_1.
2. If δ is a weighted degree in (x, y)-coordinates, then ψ is zero (see example 2.12) and consequently $p = 1$ and $\Psi = \Psi_1 = y$.

3. $\Psi_2 \equiv 0$ iff $\operatorname{ord}(\psi)p > \delta(\Psi)$.
4. For each j, $1 \leq j \leq 2$, if Ψ_j is non-zero, then $\delta(\Psi_j) = \delta(\Psi)$.

Theorem 4.1. *The semidegree δ is finitely generated if and only if $\Psi_1 \in \mathbb{C}[x, y]$.*

Before we start giving the proof of the theorem, we introduce some terminology that will be used in the proof.

Proof of theorem 4.1. At first assume δ is finitely generated. Let $A := \mathbb{C}[x, y]$ and $f_1, \ldots, f_m \in A \setminus \mathbb{C}$ such that $(1)_1, (f_1)_{\delta(f_1)}, \ldots, (f_m)_{\delta(f_m)}$ generate A^δ as a $\mathbb{C}$-algbera.

Claim 4.12.1. *For all $f \in A \setminus \mathbb{C}$,*

$$\frac{\delta(f)}{\deg(f)} \geq \min_{1 \leq j \leq m} \frac{\delta(f_j)}{\deg(f_j)}. \tag{32}$$

Proof. At first note that the quantity in the right hand side of (32) is well defined, since for all j, $1 \leq j \leq m$, $\deg(f_j) > 0$. Let $f \in A \setminus \mathbb{C}$. Then $(f)_{\delta(f)}$ can be expressed as

$$(f)_{\delta(f)} = \sum_\alpha a_\alpha (f_1)^{\alpha_1}_{\delta(f_1)} \cdots (f_m)^{\alpha_m}_{\delta(f_m)}$$

for some $\alpha := (\alpha_1, \ldots, \alpha_m) \in \mathbb{Z}_+^m$ such that $\sum_{j=1}^m \alpha_j \delta(f_j) \leq \delta(f)$ for all α such that $a_\alpha \neq 0$. Pick α such that $d := \sum_{j=1}^m \alpha_j \deg(f_j)$ is the maximum among all α such that $a_\alpha \neq 0$. Note that $d = \deg(f_1^{\alpha_1} \cdots f_m^{\alpha_m}) \geq \deg(f) > 0$ and therefore

$$\frac{\delta(f)}{\deg(f)} \geq \frac{\alpha_1 \delta(f_1) + \cdots + \alpha_m \delta(f_m)}{\alpha_1 \deg(f_1) + \cdots + \alpha_m \deg(f_m)} \geq \min_{1 \leq j \leq m} \frac{\delta(f_j)}{\deg(f_j)},$$

as required to complete the proof of the claim. □

W.l.o.g. we may assume that $\min\{\delta(f_j)/\deg(f_j) : 1 \leq j \leq m\}$ is achieved for $j = 1$. To complete the proof of ($\Rightarrow$) direction of the theorem, it suffices to show that if $\Psi_1 \notin \mathbb{C}[x, y]$, then there exists a non-constant polynomial $f \in \mathbb{C}[x, y]$ such that $\delta(f)/\deg(f) < \delta(f_1)/\deg(f_1)$. Let the factorization of f_1 in $K^*(x)[y]$ be

$$f_1 = x^e \left(\prod_{j=1}^{m_1 - 1} (y - \phi_{1j}) \right)^{e_1} \cdots \left(\prod_{j=1}^{m_n - 1} (y - \phi_{nj}) \right)^{e_n} \tag{33}$$

where $n, e, e_1, \ldots, e_n$ are non-negative integers, $\{\phi_{i0} : 1 \leq i \leq n\}$ is a collection of mutually non-conjugate degree-wise Puiseux series, and for each i, $1 \leq i \leq n$, ϕ_{ij}'s are the conjugates of ϕ_{i0}.

Claim 4.12.2. *If $\Psi_1 \notin \mathbb{C}[x, y]$, then either $e > 0$ or there exists i, $1 \leq i \leq n$, such that $\delta(y - \phi_{ij}) > r$ for all j, $0 \leq j \leq m_i - 1$.*

Proof. □

Let d be Let d be the largest non-negative integer such that Ψ divides f_1 in $K^*(x)[y]$. Since $\Psi \notin A$, Then there exists a non-constant $g \in K^*(x)$ and a nosuch that

$\left(\prod_{j=0}^{p-1} (y - \psi_j)\right)^{e_0}$ Now assume that $\Psi_1 \in \mathbb{C}[x, y]$. We go by inductions on

□

We look at We look at the discrete valuations associated to irreducible components of X_∞. Let V be an irreducible component of X_∞ and ν be the discrete valuation on $\mathbb{C}(X)$ corresponding to V, i.e. for all $f \in \mathbb{C}(X)$, $\nu(f)$ is the order of vanishing of f along V. Since V is at infinity with respect to X, it follows that (e.g. by [2, Proposition 1]) there exists $f \in \mathbb{C}[x, y]$ such $\nu(f) < 0$, therefore at least one of $\nu(x)$ and $\nu(y)$ is negative. Replacing x by $x - \lambda y$ for a generic $\lambda \in \mathbb{C}$, we may assume that $\nu(x)$ is negative.

We now inductively define a sequence of polynomials $\{y_i : i \geq 0\}$. Let $y_0 := y$. Now assume $\{y_0, \dots, y_k\}$ have been defined. Let p_k be the smallest *positive* integer such that $p_k\nu(y_k) = q_k\nu(x)$ for some integer q_k. Let $y'_k := y_k^{p_k}/x^{q_k}$. Then $\nu(y'_k) = 0$, so that y'_k restricts to a non-zero rational function on V. If $y'_k|_V$ is *not* constant, then stop. Otherwise there exists $a_k \neq 0 \in \mathbb{C}$ such that $y'_k|_V \equiv a_k$, and define

$$y_{k+1} := \begin{cases} y_k^{p_k} - a_k x^{q_k} & \text{if } q_k \geq 0, \\ y_k^{p_k} x^{-q_k} - a_k & \text{otherwise.} \end{cases}$$

Continuing in this way, we get a possibly infinite sequence $\mathcal{S} := \{y_0, y_1, \dots\}$.

Claim 4.1.3. *$\mathcal{S}$ is be finite.*

Proof. Use [1]. □

Assume $\mathcal{S}$ stops at y_k, with $k \geq 0$. Let δ be the semidegree on $A := \mathbb{C}[x, y]$ corresponding to V, i.e. $\delta := -\nu$. Our main result is that δ is finitely generated.

Theorem 4.2. *A^δ is generated by $(1)_1, (x)_{\delta(x)}, (y_0)_{\delta(y_0)}, \dots, (y_k)_{\delta(y_k)}$.*

Proof. Let $z_0, \dots, z_k$ be new indeterminates and I be the ideal in $B := \mathbb{C}[x, z_0, \dots, z_k]$ generated by $z_j - y_j$, $0 \leq j \leq k$. Then $B/I \cong A$. Let δ' be the weighted degree on B such that $\delta'(x) = \delta(x)$, and $\delta'(z_j) = \delta(y_j)$ for $0 \leq j \leq k$. Let $\bar{\delta}$ be the degree-like function on $\mathbb{C}[x, y]$ induced by δ', i.e. for all $f \in A$,

$$\bar{\delta}(f) := \min\{\delta'(g) : g \in B,\ f - g \in I\}.$$

It follows from the definition of $\bar{\delta}$ that $A^{\bar{\delta}}$ is generated by $(1)_1, (x)_{\delta(x)}, (y_0)_{\delta(y_0)}, \dots, (y_k)_{\delta(y_k)}$. Therefore it suffices to show that $\delta = \bar{\delta}$.

In the course of this proof, whenever we say 'weighted homogeneous' for an element in B, it would mean weighted homogeneous with respect to the weightings determined by δ'. For all $g \in B$, let us write $\mathfrak{L}(g)$ for the *leading form* of g with respect to the weighting determined by δ' and let J be the *leading ideal* of I with respect to δ', i.e.

$$J := \langle \mathfrak{L}(g) : g \in I \rangle.$$

Claim 4.2.1. *J is a prime ideal of B. More precisely, for $1 \leq j \leq k$, define*

$$g_j := \begin{cases} z_{j-1}^{p_{j-1}} - a_{j-1}x^{q_{j-1}} & \textit{if } q_{j-1} \geq 0, \\ z_{j-1}^{p_{j-1}} x^{-q_{j-1}} - a_{j-1} & \textit{otherwise.} \end{cases}$$

Then J is generated by $g_1, \dots, g_k$.

Proof. For each j, $1 \leq j \leq k$, $z_j - g_j \in I$ and therefore $g_j \in J$ (since g_j is weighted homogeneous and $\delta'(g_j) > \delta'(z_j)$). Therefore J contains the ideal J' of B generated by $g_1, \ldots, g_k$. Note that J' is a prime ideal of B with $B/J' \cong \mathbb{C}[z_k, x, x^{q_0/p_0}, \ldots, x^{q_{k-1}/p_{k-1}}]$. In particular, $\dim J' = 2$. On the other hand, $\operatorname{Proj} B/J$ is isomorphic to the complement of $\mathbb{C}^2$ in $X^{\bar{\delta}}$, so that J has pure dimension 2. Therefore $J = J'$. □

Let us denote the surjection $B/I \twoheadrightarrow A$ by ϕ. Let K be the ideal of B generated by weighted homogeneous elements $g \in B$ such that $\delta'(g) > \delta(\phi(g))$. We claim that K is also prime. Indeed, pick weighted homogeneous $g_1, g_2 \in B$ such that $g_1 g_2 \in K$. Then $\delta'(g_1) + \delta'(g_2) = \delta'(g_1 g_2) > \delta(\phi(g_1 g_2)) = \delta(\phi(g_1)) + \delta(\phi(g_2))$. Therefore there is j, $1 \leq j \leq 2$ such that $\delta'(g_j) > \delta(\phi(g_j))$, or equivalently, $g'_j \in K$. Consequently, K is a prime ideal of B.

We now show that $K = J$. Indeed, for all j, $1 \leq j \leq k$, $g_j \in K$, where g_j's are as in claim 4.2.1, and therefore, $K \supseteq J$. Assume to the contrary of the claim that $K \supsetneq J$. Then the image L of K in B/J is a non-zero prime ideal. Now, claim 4.2.1 implies that

$$\begin{aligned} B/J &= \mathbb{C}[x, z_0, \ldots, z_k]/\langle z_0^{p_0} - a_0 x^{q_0}, \ldots, z_{k-1}^{p_{k-1}} - a_{k-1} x^{q_{k-1}} \rangle \\ &\cong \mathbb{C}[x, x^{q_0/p_0}, \ldots, x^{q_{k-1}/p_{k-1}}, z_k] =: R \end{aligned}$$

Let $p := \delta(x) > 0$ and define

$$R' := \begin{cases} \mathbb{C}[x^{1/p}, z_k] & \text{if } q_j \geq 0 \text{ for all } j,\ 0 \leq j < k \\ \mathbb{C}[x^{1/p}, x^{-1/p}, z_k] & \text{otherwise.} \end{cases}$$

Since $q_j/p_j = \delta(y_j)/\delta(x) = \delta(y_j)/p$ for all j, $0 \leq j \leq k-1$, it follows that $R \subseteq R'$. Now, both the surjection $B/J \twoheadrightarrow R$ and the inclusion $R \hookrightarrow R'$ are homomorphisms of graded rings (where the gradings of B, R and R' are induced by δ'; in particular, the grading of R' (and R) is determined by the weighted degree that assigns weight 1 to $x^{1/p}$ and $\delta(y_k)$ to z_k). Therefore L is a non-zero prime ideal of R which is also homogeneous (with respect to the grading on R). Since R' is integral over R, there is a homogeneous prime ideal L' of R' such that $L' \cap R = L$. But then L' is generated in R' by a (weighted) homogeneous polynomial of the form $az_k - bx^{\delta(y_k)/p}$ for some $(a, b) \neq (0, 0) \in \mathbb{C}^2$. It follows that $a^p z_k^p - b^p x^{\delta(y_k)} \in L' \cap R = L$ and consequently, $g := a^p z_k^p - b^p x^{\delta(y_k)} \in K$. Note that g is weighted homogeneous with weight $p\delta(y_k)$. By definition of K, it follows that

$$\begin{aligned} \delta(\phi(g)) < p\delta(y_k) &\Rightarrow \delta(a^p y_k^p - b^p x^{\delta(y_k)}) < \delta(x^{\delta(y_k)}) \\ &\Rightarrow \delta\left(\frac{a^p y_k^p - b^p x^{\delta(y_k)}}{x^{\delta(y_k)}}\right) < 0 \\ &\Rightarrow \left.\left(\frac{a^p y_k^p}{x^{\delta(y_k)}} - b^p\right)\right|_U \equiv 0 \text{ for a Zariski open subset } U \text{ of } V. \end{aligned}$$

But the last assertion implies that $\frac{y_k^p}{x^{\delta(y_k)}}$ restricts to a constant function on V, which contradicts the choice of y_k. This contradiction proves that $K = J$, as claimed.

Now, let $f \in A$ and $d := \bar{\delta}(f)$. Then there exists $g \in B$ such that $g - f \in I$ and $\delta'(g) = d$. Let $g = \sum_{e \leq d} g_e$ be the decomposition of g into weighted homogeneous components. It follows from the definition of $\bar{\delta}$ and J that $g_d = \mathfrak{L}(g) \notin J$. Then $g_d \notin K$ (because $K = J$) and therefore $\delta(\phi(g_d)) = \delta'(g_d) = d$. On the other hand, for all $e < d$, $\delta(\phi(g_e)) \leq \delta'(g_e) \leq e < d$. It follows that $\delta(f) = \delta(\phi(g_d) + \sum_{e<d} \phi(g_e)) = \delta(\phi(g_d)) = d = \bar{\delta}(f)$. Therefore, $\delta = \bar{\delta}$, and as remarked earlier, this completes the proof of the theorem. $\square$

Definition 4.3. Let δ be a semidegree on $A := \mathbb{C}[x, y]$. We say that δ is a *divisorial semidegree* if δ is the semidegree on A associated to a component at infinity of some normal algebraic surface $\tilde{X}$ containing $\mathbb{C}^2$ as a Zariski open subset. This definition agrees with the usual notion of *divisorial valuations* (e.g. see [1]), since the negative of a divisorial semidegree is indeed a divisorial valuation.

Corollary 4.4. *Let $\bar{X}$ be a primitive normal compactification of $\mathbb{C}^2$. Then $\bar{X}$ is a weighted complete intersection. More precisely, there exist a non-negative integer k and positive integers $d, e_0, \ldots, e_k$, $p_0, \ldots, p_{k-1}$, $q_0, \ldots, q_{k-1}$, such that $p_j e_j = q_j d > e_{j+1}$ for all j, $0 \leq j \leq k-1$ and $\bar{X}$ is isomorphic to a subvariety V of the weighted projective space $\mathbf{W}\mathbb{P} := \mathbb{P}^{k+2}(1, d, e_0, \ldots, e_k)$. If the weighted homogeneous coordinates of $\mathbf{W}\mathbb{P}$ are $[w : x : z_0 : \cdots : z_k]$ corresponding respectively to weights $1, d, e_0, \ldots, e_k$, then V is defined by k (weighted homogeneous) equations $w^{r_j} z_j = z_{j-1}^{p_{j-1}} - a_{j-1} x^{q_{j-1}}$ where $r_j := q_{j-1} d - e_j$ and a_{j-1} is a non-zero complex number for $1 \leq j \leq k$.*

Corollary 4.5. *Let $\bar{X}$ be a primitive normal compactification of $\mathbb{C}^2$ and $X_\infty := \bar{X} \setminus \mathbb{C}^2$.*

1. *X_∞ is a non-singular curve isomorphic to $\mathbb{P}^1$.*
2. *X has at most two singular points.*
3. *If $\bar{X}$ is non-singular, then $\bar{X} \cong \mathbb{P}^2(\mathbb{C})$.*

Question 4.6. What does the 'curve' has one point at infinity imply in this case?

Remark 4.7. Does [5, Proposition 6.7] imply that if A^δ is not finitely generated for a semidegree δ, then there does *not* exist an algebraic surface $\bar{X}$ containing X such that $X_\infty := \bar{X} \setminus X$ is irreducible and the semidegree corresponding to X_∞ is δ? Well, we need to show that X_∞ is $\mathbb{Q}$-Cartier. Check out the remark from the second last sentence of page 1 of the other Schröer paper.

Remark 4.8. Also, can you answer question 5.10 of the other Schröer paper by taking $\delta := \max\{\delta_1, \delta_2\}$ such that both δ_j's are positive, ψ_1 has positive exponents and δ_2 is not complete? The goal would be to ensure $K_{X^\delta} \sim D_2$. It would also require to answer the question in the preceding remark in AFFIRMATIVE.

Corollary 4.9. *Let $\psi \in \mathbb{C}\langle\langle x \rangle\rangle$, $r \in \mathbb{R}$, and ψ_r be the truncation of ψ at r, i.e. $\psi_r := \sum_{\alpha > r} c_\alpha(\psi) x^\alpha$. Assume ψ_r has the form*

$$\psi_r = \left(a_1 x^{\frac{q_1}{p_1}} + \sum_{j=1}^{k_1} a_{1j} x^{\frac{q_{1j}}{p_1}} \right) + \cdots + \left(a_l x^{\frac{q_l}{p_1 p_2 \cdots p_l}} + \sum_{j=1}^{k_l} a_{l,j} x^{\frac{q_{l,j}}{p_1 p_2 \cdots p_l}} \right)$$

where $(p_1, q_1), \ldots, (p_l, q_l)$ are degree-wise Puiseux pairs of ψ_r. For each i, j, k, such that $0 \leq k \leq l$, $k + 1 \leq j$ Then there exists a polynomials $f \in \mathbb{C}[x, y]$ such that $f = 0$ has one place at infinity and the degree-wise Puiseux series associated to f has the form $y = \phi(x)$ such that $\phi \equiv_r \psi$ iff

5 Proof of the main result

Let x, y be indeterminates. We say that a polynomial $f \in \mathbb{C}[x, y]$ is *regular* in y if $\deg_y(f(0, y)) = \deg(f)$. For a $\mathbb{C}$-algebra R, we write

$$\begin{aligned} R\langle\langle x\rangle\rangle &:= \text{the ring of degree-wise Puiseux series in } x \text{ with coefficients in } R \\ R\{\{x\}\} &:= \{\phi \in R\langle\langle x\rangle\rangle : \omega \in \mathbb{Z} \text{ for all } \omega \text{ such that } c_\omega(\phi) \neq 0\}. \end{aligned}$$

Lemma 5.1. *Let $y_0, y_1, \ldots, y_l$ be indeterminates,*

$$\begin{aligned} A &:= \mathbb{C}\{\{x\}\}[y] \subseteq \mathbb{C}\langle\langle x\rangle\rangle[y] =: A^*, \\ B &:= \mathbb{C}\{\{x\}\}[y_0, y_1, \ldots, y_l] \subseteq \mathbb{C}\langle\langle x\rangle\rangle[y_0, y_1, \ldots, y_l] =: B^*, \end{aligned}$$

and $p_0, \ldots, p_l$ be positive integers. Let $\chi : B^* \to A^*$ be a $\mathbb{C}$-algebra homomorphism which maps $x \mapsto x$ and $y_k \mapsto f_k \in \mathbb{C}[x, x^{-1}, y]$ for all k, $0 \leq k \leq l$. Assume that for all k, $0 \leq k \leq l$ $d_k := \deg(f_k) = p_0p_1 \cdots p_k$ and f_k has the form

$$f_k = y^{d_k} + \sum_{j=1}^{d_k} h_{k,j}(x)y^{d_k - j}$$

with $h_{k,j} \in \mathbb{C}[x, x^{-1}]$. Let $H = \sum h_\beta(x)y^\beta \in B^*$ such that if $h_\beta(x) \neq 0$ for some $\beta := (\beta_0, \ldots, \beta_l) \in \mathbb{Z}^{l+1}_{\geq 0}$, then

$$\beta_k < p_{k+1} \text{ for all } k, 0 \leq k < l. \tag{34}$$

Then

1. *If $H \neq 0$, then $\chi(H) \neq 0$.*
2. *If $\chi(H) \in A$, then $H \in B$.*

Proof. At first note that for $\beta \in \mathbb{Z}^{l+1}_{\geq 0}$, (34) is equivalent to the following property:

$$\sum_{j=0}^{k} \beta_j d_j < d_{k+1} \text{ for all } k,\ 0 \leq k < l. \tag{34'}$$

Indeed, pick $\beta \in \mathbb{Z}^{l+1}_{\geq 0}$. It is clear that if β satisfies (34′), then it satisfies (34). So assume β satisfies (34). Then we show by induction on k that (34′) is also satisfied. Indeed, since $\beta_0 < p_1$, it follows that $\beta_0 d_0 < p_1 d_0 = p_1 p_0 = d_1$. Therefore β satisfies (34′) for $k = 0$. Assume now that β satisfies (34′) for $k - 1$ for some $k < l$. Then

$$\begin{aligned} \sum_{j=0}^{k} \beta_j d_j &= \sum_{j=0}^{k-1} \beta_j d_j + \beta_k d_k \\ &< (\beta_k + 1)d_k \qquad \text{(by induction)} \\ &\leq p_{k+1} d_k \qquad \text{(since } \beta_k < p_{k+1}\text{)} \\ &= d_{k+1}. \end{aligned}$$

It follows by induction that β satisfies ($34'$) for all k, $0 \leq k < l$. Therefore (34) and ($34'$) are indeed equivalent.

Pick $H = \sum h_\beta(x) y^\beta \in B^*$ such that for all $\beta \in \mathbb{Z}_{\geq 0}^{l+1}$, if $h_\beta(x) \neq 0$, then β satisfies (34), and therefore also ($34'$). Let ω be the weighted degree on B^* such that $\omega(x) := 0$ and $\omega(y_k) := d_k$, $0 \leq k \leq l$. Let $H = H_1 + \cdots + H_s$ be the decomposition of F into weighted homogeneous forms with $\omega(H_1) > \cdots > \omega(H_s)$.

Claim 5.1.1. *For each k, $1 \leq k \leq s$, $H_k = h_k(x) y^{\beta^k}$ for some $\beta^k \in \mathbb{Z}_{\geq 0}^{l+1}$ and $h_k(x) \in \mathbb{C}\langle\langle x \rangle\rangle$.*

Proof. Write $H_k = \sum_\beta h_{k,\beta}(x) y^\beta$. Pick β, β' such that $h_{k,\beta'}(x)$ and $h_{k,\beta'}(x)$ are non-zero. It suffices to show that $\beta = \beta'$.

Let i be the 'right-most' coordinate such that $\beta_i \neq \beta'_i$. Assume w.l.o.g. $\beta_i \geq \beta'_i$. Since H_k is weighted homogeneous with respect to ω, it follows that $d_0\beta_0 + \cdots + d_i\beta_i = d_0\beta'_0 + \cdots + d_i\beta'_i < d_i(\beta'_i + 1)$ (where the last inequality follows from ($34'$)), and therefore $\beta'_i \leq \beta_i < \beta'_i + 1$. It follows that $\beta'_i = \beta_i$. Similarly, it is straightforward to see by induction that $\beta_{i'} = \beta'_{i'}$ for all $i' \leq i$. It follows that $\beta' = \beta$ and the claim is proved. $\square$

Assume $H \neq 0$. Then claim 5.1.1 implies that $H = \sum_{k=1}^{s} h_k(x) y^{\beta^k}$ with $w_1 := \omega(y^{\beta^1}) > \cdots > \omega(y^{\beta^s})$ and $h_k(x) \neq 0$ for all k, $1 \leq k \leq s$. Then the coefficient of y^{w_1} in $\chi(H)$ is $h_1(x)$. In particular, $\chi(H) \neq 0$. This proves assertion 1 of the lemma.

Now assume $\chi(H) \in A$. If $h_k(x) \in \mathbb{C}\{\{x\}\}$ for all k, $1 \leq k \leq s$, then $H \in B$ and there is nothing to prove. So assume that there exists k such that $h_k(x) \notin \mathbb{C}\{\{x\}\}$. Let k_0 be the smallest value of k as in the preceding sentence and $H^* := H - \sum_{j<k_0} h_j(x) y^{\beta^j}$. Then $\chi(H^*) \in A$. But, if $w_0 := \omega(y^{\beta^{k_0}})$, then the coefficient of y^{w_0} in $\chi(H^*)$ is precisely $h_{k_0}(x)$, which is *not* an element of $\mathbb{C}\{\{x\}\}$. This contradiction shows that $H \in B$ and proves assertion 2 of the lemma. $\square$

Corollary 5.2. *Let $p_1, \ldots, p_i$ be positive integers and $f_1, \ldots, f_i \in \mathbb{C}[x, x^{-1}, y]$ such that for all k, $1 \leq k \leq i$, $d_k := \deg(f_k) = p_1 \cdots p_k$ and f_k has the form*

$$f_k = y^{d_k} + \sum_{j=1}^{d_k} h_{k,j}(x) y^{d_k - j}$$

with $h_{k,j} \in \mathbb{C}[x, x^{-1}]$. Set $f_0 := y$ and $B_i := \mathbb{C}\langle\langle x \rangle\rangle[y_0, \ldots, y_i]$. Let $\Phi \in \mathbb{C}\langle\langle x \rangle\rangle[y]$ and

$$p_{i+1} := \left\lceil \frac{\deg_y(\Phi)}{p_1 p_2 \cdots p_i} \right\rceil = \min\{k : k p_1 \ldots p_i \geq \deg_y(\Phi)\}.$$

1. *If $p_1 p_2 \cdots p_i$ does not divide $\deg_y(\Phi)$, or equivalently $\deg_y(\Phi) < p_1 p_2 \cdots p_{i+1}$, then there is a unique $F_\Phi \in B_i$ such that $\Phi = F_\Phi(x, f_0, \ldots, f_i)$ and $\deg_{y_k}(F_\Phi) < p_{k+1}$ for all k, $0 \leq k \leq i$.*

2. *If $p_1 p_2 \cdots p_i$ divides $\deg_y(\Phi)$, or equivalently $\deg_y(\Phi) = p_1 p_2 \cdots p_{i+1}$, then there is a unique $F_\Phi \in B_i$ such that $\Phi = F_\Phi(x, f_0, \ldots, f_i)$ and $\deg_{y_k}(F_\Phi - y_i^{p_{i+1}}) < p_{k+1}$ for all k, $0 \leq k \leq i$.*

Moreover, if F_Φ is as in assertion 1 or 2 above, then it satisfies following additional properties:

3. *The polydromy order of F_Φ equals the polydromy order of Φ. If $\Phi \in \mathbb{C}[x, x^{-1}, y]$, then $F_\phi \in \mathbb{C}[x, x^{-1}, y_0, \dots, y_i]$.*
4. *Let ω be the weighted degree on B_i such that $\omega(x) := 0$, $\omega(y_0) := 1$ and $\omega(y_k) := d_k$, $1 \leq k \leq i$. Then $\omega(F_\Phi) = \deg_y(\Phi)$.*

Proof. At first note that the uniqueness of F_Φ as in the lemma is a straightforward application of lemma 5.1. Indeed, Indeed, let $F_1, F_2 \in \mathbb{C}\langle\langle x \rangle\rangle[y_0, \dots, y_{i-1}]$ such that for each n, $1 \leq n \leq 2$, $F_n(x, f_0, \dots, f_{i-1}) = \Phi^{(i)}$ and $\deg_{y_k}(F_n - y_{i-1}^{p_i}) < p_{k+1}$ for all k, $0 \leq k < i$. Then $(F_1 - F_2)(x, f_0, \dots, f_{i-1}) = 0$. Applying lemma 5.1 (with l replaced by $i-1$ and $p_0 = 1$) to $F_1 - F_2$, it follows that $F_1 = F_2$. Therefore, if there exists F_i^ϕ as in the lemma, then it is unique.

Now we prove the existence part of the lemma. Let $G_0 := \Phi^{(i)}(x, y_0) \in \mathbb{C}\langle\langle x \rangle\rangle[y_0] \subseteq B_{i-1}$. Then G_0 is monic in y_0 of degree $p_1 \cdots p_i$. We now construct a sequence $\{G_n : n \geq 0\}$ of the elements of B_{i-1} as follows:

Assume G_n is given. Write $G_n := \sum g_{n,\beta}(x) y^\beta$. For every $\beta := (\beta_0, \dots, \beta_{i-1})$ such that $g_{n,\beta}(x)$ is non-zero, let

$$k_\beta := \begin{cases} 0 & \text{if } \beta_0 < p_1, \\ \max\{k \leq i-1 : p_1 p_2 \cdots p_k \leq \beta_0\} & \text{otherwise.} \end{cases}$$

$$d_\beta := \begin{cases} 0 & \text{if } k_\beta = 0, \\ \max\{d : d p_1 p_2 \cdots p_{k_\beta} \leq \beta_0\} & \text{otherwise.} \end{cases}$$

$$\hat{g}_{n,\beta} := y_0^{\beta_0 - d_\beta p_1 \cdots p_{k_\beta}} y_1^{\beta_1} \cdots y_{i-1}^{\beta_{i-1}} \left(y_{k_\beta} - \left(f_{k_\beta}(x, y_0) - y_0^{p_1 \cdots p_{k_\beta}} \right) \right)^{d_\beta}.$$

Note that if $d_\beta = 0$, then $\hat{g}_{n,\beta} = y^\beta$. Let

$$G_{n+1} := \sum_\beta g_{n,\beta}(x) \hat{g}_{n,\beta}(y_0, \dots, y_{i-1}).$$

Inductive assumption $5_{(k)}$ implies that

$$\deg_{y_0} \left(f_{k_\beta}(x, y_0) - y_0^{p_1 \cdots p_{k_\beta}} \right) < p_1 \cdots p_{k_\beta}$$

for all β and therefore $\deg_{y_0}(G_{n+1}) \leq \deg_{y_0}(G_n)$ for every n, and therefore there exists n_0 such that $\deg_{y_0}(G_n) = \deg_{y_0}(G_{n_0})$ for all $n \geq n_0$. Then it follows from the construction of G_n's that $\deg_{y_0}(G_{n_0}) < p_1$ and $G_n = G_{n_0}$ for all $n \geq n_0$. Let $F_i^\phi := G_{n_0}$. We claim that F_i^ϕ satisfies the conditions of assertion **??** of the lemma. Indeed, note from the definition of $\hat{g}_{n,\beta}$ that

$$\hat{g}_{n,\beta}|_{y_{k_\beta} = f_{k_\beta}(x, y_0)} = y^\beta|_{y_{k_\beta} = f_{k_\beta}(x, y_0)},$$

and therefore, $G_{n+1}(x, f_0, \dots, f_{i-1}) = G_n(x, f_0, \dots, f_{i-1})$ for all $n \geq 0$. In particular,

$$F_i^\phi(x, f_0, \dots, f_{i-1}) = G_{n_0}(x, f_0, \dots, f_{i-1}) = G_0(x, y) = \Phi^{(i)}.$$

Now note that $\deg_{y_0}(G_0) = p_1 \cdots p_i$ and $y_0^{p_1 \cdots p_i}$ is a monomial in G_0. It follows from the construction of G_1 that $y_{i-1}^{p_i}$ is a monomial in G_1, $G_1 \in \mathbb{C}\langle\langle x \rangle\rangle[y_0, y_{i-1}]$, and if the coefficient of $y_0^{\beta_0} y_{i-1}^{\beta_{i-1}}$ in $G_1 - y_{i-1}^{p_i}$ is non-zero, then

$$\beta_0 + p_1 \cdots p_{i-1}\beta_{i-1} < p_1 \cdots p_i \quad \Rightarrow \quad \frac{\beta_0}{p_1 \cdots p_{i-1}\beta_{i-1}} + \beta_{i-1} < p_1.$$

It follows from the construction process of G_n's that for all $n \geq 0$, $\deg_{y_{i-1}}(G_n - y_{i-1}^{p_i})$ is bounded by

$$\max\left\{\frac{\beta_0}{p_1 \cdots p_{i-1}\beta_{i-1}} + \beta_{i-1} : \text{coefficient of } y_0^{\beta_0} y_{i-1}^{\beta_{i-1}} \text{ in } G_1 - y_{i-1}^{p_i} \text{ is non-zero}\right\} < p_1.$$

A straightforward induction on $i - k$ now shows that $\deg_{y_k}(G_n - y_{i-1}^{p_i}) < p_{k+1}$ for all k, $0 \leq k < i$ and for all $n \geq 0$. This completes the proof of the lemma. □

Lemma 5.3. *Let $y_0, y_1, \ldots, y_l$ be indeterminates and $B^* := \mathbb{C}\langle\langle x \rangle\rangle[y_0, y_1, \ldots, y_l]$. Let $\pi : B^* \to A^* := \mathbb{C}\langle\langle x \rangle\rangle[y_0]$ be a $\mathbb{C}$-algebra homomorphism that sends $x \mapsto x$, $y_0 \mapsto y_0$ and $y_j \to f_j \in \mathbb{C}[x, x^{-1}, y_0]$ for $1 \leq j \leq l$. For all $\Phi \in B^*$, write $\Phi = \sum_\alpha c_\alpha^\Phi(y) x^\alpha$. Recall that the polydromy order of $\Phi \in B^*$ is the smallest positive integer s such that $\Phi \in \mathbb{C}\{\{x^{\frac{1}{s}}\}\}[y_0, y_1, \ldots, y_l]$. If $\Phi \in B^*$ and s is a multiple of the polydromy order of Φ and μ is an s-th root of unity, then define*

$$\mu \star_s \Phi := \sum_i c_{i/s}^\Phi(y_0, \ldots, y_l) \mu^i x^{i/s}.$$

The claim is: $\pi(\mu \star_s \Phi) = \mu \star_s \pi(\Phi)$.

Proof. Let Φ, s and μ be as in the statement of the lemma. Note that

$$\begin{aligned}
\pi(\mu \star_s \Phi) &= \sum_i c_{i/s}^\Phi(f_0, \ldots, f_l)\mu^i x^{i/s}, \text{ and} \\
\mu \star_s \pi(\Phi) &= \mu \star_s \left(\sum_i c_{i/s}^\Phi(y_0, f_1, \ldots, f_l) x^{i/s}\right) \\
&= \sum_i \mu \star_s \left(c_{i/s}^\Phi(y_0, f_1, \ldots, f_l) x^{i/s}\right).
\end{aligned}$$

Since $f_i \in \mathbb{C}[x, x^{-1}, y_0]$ for $1 \leq i \leq l$, it follows that $c_{i/s}^\Phi(y_0, f_1, \ldots, f_l)x^{i/s} \in \mathbb{C}[x, x^{-1}, y_0]$ for all $i \in \mathbb{Z}$. Now, we claim that if $g \in \mathbb{C}[x, x^{-1}, y_0]$, then $\mu \star_s (g x^{\frac{i}{s}}) = \mu^i g x^{\frac{i}{s}}$. Indeed, let $g = \sum a_{j,k} x^j y_0^k$. Then

$$\mu \star_s (g x^{\frac{i}{s}}) = \mu \star_s \left(\sum a_{j,k} x^{\frac{i+js}{s}} y_0^k\right) = \sum a_{j,k} \mu^{i+js} x^{\frac{i+js}{s}} y_0^k = \mu^i \sum a_{j,k} x^{\frac{i+js}{s}} y_0^k = \mu^i g x^{\frac{i}{s}}$$

as claimed. It follows that $\mu \star_s \left(c_{i/s}^\Phi(y_0, f_1, \ldots, f_l) x^{i/s}\right) = \mu^i c_{i/s}^\Phi(y_0, f_1, \ldots, f_l) x^{i/s}$ for all i, and consequently, $\mu \star_s \pi(\Phi) = \pi(\mu \star_s \Phi)$, as required. □

Corollary 5.4. *Let $\phi \in \mathbb{C}\langle\langle x \rangle\rangle$ with Puiseux pairs $(p_1, q_1), \ldots, (p_l, q_l)$. Let i be an integer with $1 \le i \le l$ and $f_1, \ldots, f_{i-1} \in \mathbb{C}[x, x^{-1}, y]$ such that for all k, $1 \le k \le i-1$, $d_k := \deg(f_k) = p_1 \cdots p_k$ and f_k has the form*

$$f_k = y^{d_k} + \sum_{j=1}^{d_k} h_{k,j}(x) y^{d_k - j}$$

with $h_{k,j} \in \mathbb{C}[x, x^{-1}]$. Set $f_0 := y$. Then there is a unique $F_i^{\phi} \in \mathbb{C}\langle\langle x \rangle\rangle[y_0, \ldots, y_{i-1}] =: B_{i-1}$ such that $\Phi^{(i)} = F_i^{\phi}(x, f_0, \ldots, f_{i-1})$ and $\deg_{y_k}(F_i^{\phi} - y_{i-1}^{p_i}) < p_{k+1}$ for all k, $0 \le k < i$. If $\phi \in \mathbb{C}[x, x^{-1}, y]$, then $F_i^{\phi} \in \mathbb{C}[x, x^{-1}, y_0, \ldots, y_{i-1}]$. $\square$

Let ψ be a *finite* degree-wise Puiseux series in x with Puiseux pairs $(p_1, q_1), \ldots, (p_l, q_l)$ and $r \in \mathbb{Q}$ with $r < \operatorname{ord}_x(\psi)$. Define

$$r_k := \begin{cases} \frac{q_k}{p_1 p_2 \cdots p_k} & \text{if } 1 \le k \le l, \\ r & \text{if } k = l+1, \end{cases}$$

For each k, $0 \le k \le l$, let ξ_k be an indeterminate, ζ_k be a $(p_1 \ldots p_k)$-th root of unity (for $k = 0$, we define $\zeta_0 := 1$ and $p_0 := 1$), and

$$\psi_k(x) := \sum_{\omega > r_{k+1}} c_\omega(\psi) x^\omega, \qquad \tilde{\psi}_k(x, \xi_k) := \psi_k(x) + \xi_k x^{r_{k+1}},$$

$$\Psi_k^*(x, y) := \prod_{j=1}^{p_0 \cdots p_k} (y - \zeta_k^j \star_{p_0 \cdots p_k} \psi_k(x)),$$

$$\delta_k := \text{the semidegree on } \mathbb{C}[x, y] \text{ defined by}$$

$$\delta_k(f) := \deg_x(f(x, \tilde{\psi}_k(x, \xi_k))) \text{ for all } f \in \mathbb{C}[x, y].$$

Below we determine $\delta_k(\Psi_j^*)$ for $0 \le j \le k \le l$. For each pair of positive integers m, n such that $1 \le n \le m \le l+1$, let

$$\hat{\sigma}_{m,n} := \sum_{1 \le i_1 < \ldots < i_n \le m} (p_{i_n} - 1)(p_{i_{n-1}} - 1) \cdots (p_{i_1} - 1) r_{i_1},$$

and for all k, $0 \le k \le l$, define

$$\omega_k := \sum_{j=1}^{k} \hat{\sigma}_{k,j} + r_{k+1}.$$

Lemma 5.5. *For all j, $0 \le j \le l$,*

1. $\deg_{\mathrm{x}}^{\mathrm{inv}} \left(\Psi_j^*(x, \tilde{\psi}_j(x, \xi_j)) \right) = \delta_j \left(\Psi_j^* \right) = \omega_j$, *and*
2. $\delta_k \left(\Psi_j^* \right) = \omega_j$ *for all k, $j \le k \le l$;*
3. *if $\phi \equiv_{r_{j+1}} \psi$, then assertion 1 remains true with Ψ_j^* replaced by $\Phi^{(j)}$.*

Proof. Note that for all j, k such that $0 \leq j \leq k \leq l$ and for all i, $0 \leq i < p_1 p_2 \cdots p_j$,

$$\deg_x(\tilde{\psi}_k(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x)) = \deg_x(\tilde{\psi}_j(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x))$$

and therefore

$$\begin{aligned}
\delta_k\left(\Psi_j^*\right) &= \sum_{i=0}^{p_0 \cdots p_j - 1} \deg_x(\tilde{\psi}_k(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x)) \\
&= \sum_{i=0}^{p_0 \cdots p_j - 1} \deg_x(\tilde{\psi}_j(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x)) \\
&= \delta_j\left(\Psi_j^*\right),
\end{aligned}$$

which proves assertion 2 of the lemma.

We prove assertion 1 of the lemma by induction on j. For $j = 0$, assertion 1 follows directly from definitions of δ_0 and Ψ_0^*. So assume that the assertion holds for $j-1$ for $1 \leq j \leq l$. We show that it continues to hold for j.

Let $\hat{\psi}_j := \psi_j - \psi_{j-1}$. Then

$$\begin{aligned}
\Psi_j^* &= \prod_{i=0}^{p_j - 1} \prod_{k=0}^{p_0 \cdots p_{j-1} - 1} (y - \zeta_j^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_j} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \\
&= \prod_{i=0}^{p_j - 1} \prod_{k=0}^{p_0 \cdots p_{j-1} - 1} (y - \zeta_{j-1}^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \\
&= \prod_{i=0}^{p_j - 1} \prod_{k=0}^{p_0 \cdots p_{j-1} - 1} (y - \zeta_{j-1}^{k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \\
&= \prod_{i=0}^{p_j - 1} (y - \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1}} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \\
&\quad \times \prod_{i=0}^{p_j - 1} \prod_{k=1}^{p_0 \cdots p_{j-1} - 1} (y - \zeta_{j-1}^{k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1} + k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x))
\end{aligned}$$

Since the leading term of $\hat{\psi}_j(x)$ is $a_j x^{\omega_{0,j}}$, it follows that

$$\begin{aligned}
&\delta_2\left(\prod_{i=0}^{p_j - 1} (y - \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1}} \star_{p_0 \cdots p_j} \hat{\psi}_j(x))\right) \\
&\qquad = \deg_x\left(\prod_{i=0}^{p_j - 1} (\tilde{\psi}_j(x) - \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1}} \star_{p_0 \cdots p_j} \hat{\psi}_j(x))\right) \\
&\qquad = (p_j - 1) r_j + r_{j+1}.
\end{aligned}$$

On the other hand, for each i, $0 \leq i \leq p_j - 1$,

$$\begin{aligned}
&\delta_2 \left(\prod_{k=1}^{p_0 \cdots p_{j-1}-1} (y - \zeta_{j-1}^{k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1}+k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \right) \\
&\quad = \deg_x \left(\prod_{k=1}^{p_0 \cdots p_{j-1}-1} (\tilde{\psi}_j(x) - \zeta_{j-1}^{k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x) - \zeta_j^{ip_0 \cdots p_{j-1}+k} \star_{p_0 \cdots p_j} \hat{\psi}_j(x)) \right) \\
&\quad = \deg_x \left(\prod_{k=1}^{p_0 \cdots p_{j-1}-1} (\tilde{\psi}_{j-1}(x) - \zeta_{j-1}^{k} \star_{p_0 \cdots p_{j-1}} \psi_{j-1}(x)) \right) \\
&\quad = \deg_x \left(\frac{\Psi_{j-1}^{*}(x, \tilde{\psi}_{j-1}(x, \xi_{j-1}))}{\tilde{\psi}_{j-1}(x) - \psi_{j-1}(x)} \right) \\
&\quad = \delta_{j-1} \left(\Psi_{j-1}^{*} \right) - r_j \\
&\quad = \omega_{j-1} - r_j, \quad \text{(by induction)} \\
&\quad = \sum_{k=1}^{j-1} \hat{\sigma}_{j-1,k}.
\end{aligned}$$

It follows that

$$\begin{aligned}
\delta_j(\Psi_j^{*}) &= (p_j - 1) r_j + r_{j+1} + p_j \sum_{k=1}^{j-1} \hat{\sigma}_{j-1,k} \\
&= \sum_{k=1}^{j-1} (p_j - 1) \hat{\sigma}_{j-1,k} + \sum_{k=1}^{j-1} \hat{\sigma}_{j-1,k} + (p_j - 1) r_j + r_{j+1} \\
&= (p_j - 1) \hat{\sigma}_{j-1,j-1} + \sum_{k=2}^{j-1} ((p_j - 1) \hat{\sigma}_{j-1,k-1} + \hat{\sigma}_{j-1,k}) + (\hat{\sigma}_{j-1,1} + (p_j - 1) r_j) + r_{j+1} \\
&= \sum_{k=1}^{j} \hat{\sigma}_{j,k} + r_{j+1} \\
&= \omega_j.
\end{aligned}$$

Moreover, the leading term of $\Psi_j^{*}(x, \tilde{\psi}_j(x, \xi_j))$ is $c \xi_j x^{\omega_j}$ for some non-zero $c \in \mathbb{C}$. Therefore $\deg_{\mathrm{x}}^{\mathrm{inv}}(\Psi_j^{*}(x, \tilde{\psi}_j(x, \xi_j))) = \omega_j$ as well. This completes the proof of assertion 1 of the lemma by induction.

For assertion 3, fix j, $0 \leq j \leq l$ and let $\phi \in \mathbb{C}\langle\langle x \rangle\rangle$ such that $\phi \equiv_{r_{j+1}} \psi$. Then the first j Puiseux pairs of ϕ are (p_i, q_i), $1 \leq i \leq j$. Therefore $p_1 p_2 \cdots p_j$ divides the polydromy order s of ϕ, and if $\hat{\phi} := \phi - \psi_j$, the $\deg(\hat{\phi}) \leq r_{j+1}$. Let μ be an s-th root of identity. It follows that for all i, $0 \leq i < p_1 p_2 \cdots p_j$,

$$\tilde{\psi}_j(x) - \mu^i \star_s \phi(x) = \tilde{\psi}_j(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x) - \mu^i \star_s \hat{\phi}(x).$$

In particular, for $0 < i < p_1 p_2 \cdots p_j$, the leading terms of $\tilde{\psi}_j(x) - \mu^i \star_s \phi(x)$ and $\tilde{\psi}_j(x) - \zeta_j^i \star_{p_0 \cdots p_j} \psi_j(x)$ are the same and of the form $c_i x^{\omega_i}$ for some $c_i \in \mathbb{C}$ and $\omega_i \in \mathbb{Q}$. On the other hand, for $i = 0$, the leading term of $\tilde{\psi}_j(x) - \psi_j(x)$ is $\xi_j x^{r_{j+1}}$ and the leading term of $\tilde{\psi}_j(x) - \phi(x)$ is $(\xi_j + b)x^{r_{j+1}}$ for some $b \in \mathbb{C}$. It follows that

$$\begin{aligned}\deg_{\mathrm{x}}^{\mathrm{inv}}\left(\Psi_j^*(x, \tilde{\psi}_j(x, \xi_j))\right) &= \deg_x\left(\Psi_j^*(x, \tilde{\psi}_j(x, \xi_j))\right)\\ &= \deg_x\left(\Phi^{(j)}(x, \tilde{\psi}_j(x, \xi_j))\right) = \deg_{\mathrm{x}}^{\mathrm{inv}}\left(\Phi^{(j)}(x, \tilde{\psi}_j(x, \xi_j))\right),\end{aligned}$$

which proves assertion 3 and completes the proof of the lemma. □

Lemma 5.6. *1. For all k, $0 \le k < l$, $\omega_{k+1} = p_{k+1}\omega_k - (r_{k+1} - r_{k+2})$. In particular, $\omega_{k+1} < p_{k+1}\omega_k$ for $0 \le k < l$.*

2. For all $s \in \frac{1}{p_1 p_2 \cdots p_l}\mathbb{Z}$, there are unique integers $\alpha, \beta_0, \ldots, \beta_{l-1}$ such that $\alpha + \sum_{k=0}^{l-1}\beta_k\omega_k = s$ and $0 \le \beta_k < p_{k+1}$ for all k, $0 \le k \le l-1$. Moreover, if $s \in \frac{1}{p_1 p_2 \cdots p_k}\mathbb{Z}$ for some $k \le l$, then $\beta_j = 0$ for all $j \ge k$.

Proof. Note that for $0 \le k < l$,

$$\begin{aligned}p_{k+1}\omega_k &= p_{k+1}\sum_{j=1}^{k}\hat{\sigma}_{k,j} + p_{k+1}r_{k+1}\\ &= \sum_{j=1}^{k}(p_{k+1}-1)\hat{\sigma}_{k,j} + \sum_{j=1}^{k}\hat{\sigma}_{k,j} + (p_{k+1}-1)r_{k+1} + r_{k+1}\\ &= (p_{k+1}-1)\hat{\sigma}_{k,k} + \sum_{j=2}^{k}((p_{k+1}-1)\hat{\sigma}_{k,j-1} + \hat{\sigma}_{k,j}) + (\hat{\sigma}_{k,1} + (p_{k+1}-1)r_{k+1}) + r_{k+1}\\ &= \sum_{j=1}^{k+1}\hat{\sigma}_{k+1,j} + r_{k+1}\\ &= \omega_{k+1} + (r_{k+1} - r_{k+2})\end{aligned} \tag{35}$$

This proves assertion 1 of the lemma. For the second assertion, set $\omega_{-1} := 1$ and $r_0 := 1$, and fix k, $0 \le k \le l$. For $s_1, \ldots, s_m \in \mathbb{Q}$ let us write $\langle s_1, \ldots, s_m\rangle$ for the group generated by $s_1, \ldots, s_m$ under addition and subtraction. It follows from the definition of ω_j's that

$$S_{k-1} := \langle\omega_{-1}, \omega_0, \ldots, \omega_{k-1}\rangle = \langle r_0, r_1, \ldots, r_k\rangle = \langle 1, \frac{q_1}{p_1}, \ldots, \frac{q_k}{p_1 \cdots p_k}\rangle = \frac{1}{p_1 \cdots p_k}\mathbb{Z}.$$

Let $s \in \frac{1}{p_1 \cdots p_k}\mathbb{Z}$. It follows that there are $\beta_{-1}, \ldots, \beta_{k-1} \in \mathbb{Z}$ such that $s = \sum_{j=-1}^{k-1}\beta_j\omega_j$. Moreover, note that for each j, $0 \le j \le k-1$, $p_{j+1}\omega_j \in S_{j-1}$ and therefore we may replace multiples of $p_{j+1}\omega_j$ by a linear combination of $\omega_{-1}, \ldots, \omega_{j-1}$. Doing such substitutions for $j = k-1$, and then continuing with $j = k-2, k-3, \ldots, 0$, we may assume that $0 \le \beta_j < p_{j+1}$ for $0 \le j \le j-1$.

It remains to show the uniqueness of expressions of $s = \sum_{j=-1}^{l-1}\beta_j\omega_j$ such that $0 \le \beta_j < p_{j+1}$ for $0 \le j \le l-1$. For this, it suffices to show that if $\beta := (\beta_{-1}, \ldots, \beta_{l-1}) \in \mathbb{Z}^{l+2}$ is such that $\sum_{j=-1}^{l-1}\beta_j\omega_j = 0$ and $|\beta_j| < p_{j+1}$ for $0 \le j \le l-1$, then $\beta = 0$. We give a proof of this fact by contradiction: so assume that $\beta \neq 0$. Let k be the 'right-most' coordinate of β such that $\beta_k \neq 0$. Then $k \geq 0$, and

$$\sum_{j=-1}^{k} \beta_j \omega_j = 0 \quad \iff \quad \beta_k \omega_k = -\sum_{j=0}^{k-1} \beta_j \omega_j.$$

Now, it follows from the definition of ω_j's that

$$\begin{aligned} \omega_k &= \frac{s_k}{p_1 \cdots p_k} + \frac{q_{k+1}}{p_1 \cdots p_{k+1}} \text{ for some } s_k \in \mathbb{Z}, \text{ and} \\ \omega_j &= \frac{s'_j}{p_1 \cdots p_k} \text{ for some } s'_j \in \mathbb{Z} \text{ for all } j,\ 0 \leq j < k. \end{aligned}$$

It then follows that $\beta_k(s_k p_{k+1} + q_{k+1}) = -p_{k+1} \sum_{j=-1}^{k-1} \beta_j s'_j$, and therefore p_{k+1} divides $\beta_k q_{k+1}$. Since $\gcd(p_{k+1}, q_{k+1}) = 1$, it follows that p_{k+1} divides β_k. This is a contradiction, since $|\beta_k| < p_{k+1}$. This proves that $\beta = 0$ and therefore completes the proof of the lemma. □

Theorem 5.7. *Let δ_1 and δ_2 be divisorial semidegrees on $A := \mathbb{C}[x, y]$. If δ_1 is a* positive *weighted degree, then $\delta := \max\{\delta_1, \delta_2\}$ is a finitely generated subdegree.*

Proof. At first we prove the theorem in a special case and then we reduce the general case to that special case. So assume that

H1. $\delta_1(x) = q$ and $\delta_1(y) = qr$, where q, r are positive integers,

H2. the degree-wise Puiseux series $\psi(x)$ associated to δ_2 is either identically zero or has the form

$$\psi(x) = \sum_{k=1}^{l} a_k x^{s_k}, \tag{36}$$

where s_k is an integer for each k, $1 \leq k \leq l$ and $s_1 < r$, and

H3. the generic term associated with δ_2 is ξx^s, where s is an integer with $s < s_l$.

Note that ψ is identically zero if and only if δ_2 is a weighted degree in (x, y)-coordinates. Since the maxima of weighted degrees in same coordinates are finitely generated (these are degree-like functions corresponding to toric varieties determined by convex rational polytopes, see [3, Example 3.5]), it follows that δ is finitely generated if ψ is identlically zero. So assume ψ is as in (36). If s_1 is positive, then m be the largest integer such that s_m is non-negative and let $\tilde{y} := y - \sum_{k=1}^{m} a_k x^{s_k}$. Since $s_1 < r$, it follows that δ_1 is a weighted degree in $(x, \tilde{y})$-coordinates as well, and H1 is also satisfied with y substituted by $\tilde{y}$. Therefore, replacing y by $\tilde{y}$, we may assume that the $\psi(x)$ has the form

$$\psi(x) = \sum_{k=1}^{l} a_k x^{-s_k}, \tag{36'}$$

where $\{s_k : 1 \leq k \leq l\}$ is an *increasing* sequence of *positive* integers, and the generic term associated to δ_2 is ξx^{-s}, where s is an integer with $s > s_l$. W.l.o.g. we may also assume that

$p := \delta_2(x) > q$ (for otherwise $\delta = \delta_1$ and therefore δ is finitely generated).

Let X^{δ_1} be the completion of $X := \mathbb{C}^2$ determined by δ_1. Then X^{δ_1} is a weighted projective space. We construct X^{δ} as the normalization of a 'weighted blow-up' of X^{δ_1}. Two of the components of the blow-up map are polynomials of the form $f_1 := x^{\alpha_1}y^{\beta_1}$ and $f_2 := x^{\alpha_2}(y - \psi(x))^{\beta_2}$ such that $\delta_1(f_j) = \delta_2(f_j)$ for $j = 1$ and $j = 2$. Note that

$$\delta_1(x^{\alpha_1}y^{\beta_1}) = \delta_2(x^{\alpha_1}y^{\beta_1}) \Leftrightarrow q\alpha_1 + qr\beta_1 = p\alpha_1 - ps_1\beta_1 \quad \Leftrightarrow (p-q)\alpha_1 = (qr + ps_1)\beta_1, \tag{37}$$

and for each (α_1, β_1) satisfying (37),

$$\delta_1(x^{\alpha_1}y^{\beta_1}) = q\beta_1(\frac{qr + ps_1}{p-q} + r) = pq\beta_1\frac{r + s_1}{p-q} \tag{38}$$

Similarly,

$$\begin{aligned} \delta_1(x^{\alpha_2}(y - \psi(x))^{\beta_2}) = \delta_2(x^{\alpha_2}(y - \psi(x))^{\beta_2}) &\Leftrightarrow q\alpha_2 + qr\beta_2 = p\alpha_2 - ps\beta_2 \\ &\Leftrightarrow (p-q)\alpha_2 = (qr + ps)\beta_2 \qquad (37') \end{aligned}$$

and for each (α_2, β_2) satisfying $(37')$,

$$\delta_2(x^{\alpha_2}(y - \psi(x))^{\beta_2}) = q\beta_2(\frac{qr + ps}{p-q} + r) = pq\beta_2\frac{r + s}{p-q} \tag{38'}$$

Let $\beta_1 := r(p-q)(r+s)$ and $\beta_2 := r(p-q)(r+s_1)$. Then

$$\alpha_1 = r(r+s)(qr + ps_1), \quad \alpha_2 = r(r+s_1)(qr+ps), \quad \text{and} \tag{39}$$

$$\delta_i(f_j) = pqr(r+s)(r+s_1) \text{ for all } i, j,\ 1 \leq i, j \leq 2. \tag{40}$$

Let $\Psi : \mathbb{C}^2 \to \mathbb{P}^4$ be the map defined by

$$(x, y) \mapsto (1 : y^{p(r+s)(r+s_1)} : x^{qr(r+s)(r+s_1)} : f_1 : f_2).$$

Note that $X^{\delta_1} \cong \mathbb{P}^2(1, q, qr) \cong \mathbb{P}^2(1, 1, r)$ (the latter isomorphism is a consequence of the fact that $\mathbb{P}^2(1, 1, r)$ is the image of $\mathbb{P}^2(1, q, qr)$ under the q-uple Veronese embedding, see e.g. [**?**, Section 1.3.1]). Let the weighted homogeneous coordinates of $\mathbb{P}^2(1, 1, r)$ be $(W : X : Y)$, so that $x = X/W$ and $y = Y/W^r$. Identifying X^{δ_1} with $\mathbb{P}^2(1, 1, r)$ we see that Ψ extends to a rational map (which we also denote by Ψ) from X^{δ_1} to $\mathbb{P}^4$ such that

$$\begin{aligned} \Psi(W : X : Y) = \Big(&W^{cpr} : Y^{cp} : W^{cr(p-q)}X^{cqr} : X^{\alpha_1}Y^{\beta_1} \\ &: X^{\alpha_2 - \beta_2 s_l}(X^{s_l}Y - \sum_{k=1}^{l} a_k X^{s_l - s_k}W^{r+s_k})^{\beta_2}\Big), \end{aligned}$$

where $c := (r+s)(r+s_1)$. It follows from the above expression for Ψ that there is only one point of indeterminacy of Ψ, namely the point O with weighted homogeneous coordinates $[0 : 1 : 0]$. Consequenly, the closure $\bar{\Gamma}$ of the graph Γ of $\Psi|_{X^{\delta_1}\setminus\{O\}}$ in $X^{\delta_1} \times \mathbb{P}^4$ is a disjoint union of Γ and a closed subvariety E such that E projects to P under the natural projection $\Gamma \to X^{\delta_1}$. We next show that E is an irreducible curve and find a family of curves in X^{δ_1} approaching O such that their images in Γ approach generic points of E.

Claim 5.7.1.

1. *E is an irreducible curve.*
2. *For all $b \in \mathbb{C}$, let $\psi_b(x) := \psi(x) + bx^{-s}$ and $\gamma_b(t)$ be the curve on $\mathbb{C}^2$ with associated degree-wise Puiseux series $y = \psi_b(x)$ (i.e. $\gamma_b(t) = (t, \psi_b(t))$ for all $t \in \mathbb{C} \setminus \{0\}$). Then as $t \to \infty$, $\Psi(\gamma_b(t))$ approaches a point on E.*
3. *Let $X^{\delta_1}_\infty := X^{\delta_1} \setminus X$ and P be the point of intersection of E and the closure of $\Psi(X^{\delta_1}_\infty)$ in $\bar{\Gamma}$. Then for all $z \in E \setminus \{P\}$, $z = \lim_{t\to\infty} \Psi(\gamma_b(t))$ for some $b \in \mathbb{C}$.*
4. *Conversely, for all $z \in E\setminus\{P\}$, if $\gamma(t)$ is any curve on $\mathbb{C}^2$ such that $\lim_{t\to\infty} \Psi(\gamma(t)) = z$, then there exists $b \in \mathbb{C}$ such that the degree-wise Puiseux expansion (in terms of x) for γ near z is*

   $$y = \psi_b(x) + l.o.t. \tag{41}$$

   where l.o.t. is a shortcut for 'lower order terms' (in x).
5. *Moreover, for each $b \in \mathbb{C}$, if the degree-wise Puiseux expansion for γ is as in (41), then $\lim_{t\to\infty} \Psi(\gamma(t)) = \lim_{t\to\infty} \Psi(\gamma_b(t))$.*

Proof. There is an open affine neighborhood U of O such that $U \cong \mathbb{C}^2$ with coordinates $w_0 := W/X = 1/x$ and $y_0 := Y/X^r = y/x^r$. With respect to (w_0, y_0) coordinates, Ψ has the following form:

$$\Psi(w_0, y_0) = \left({w_0}^{cpr} : {y_0}^{cp} : w_0^{cr(p-q)} : {y_0}^{\beta_1} : (y_0 - \sum_{k=1}^{l} a_k {w_0}^{r+s_k})^{\beta_2} \right), \tag{42}$$

For every $z \in E$, there is an algebraic curve $\gamma(t) \in U$ such that $\gamma(0) = O$ and $\lim_{t\to 0} \Psi(\gamma(t)) = z$. Since O has the coordinates $(0, 0)$, we may assume that near O, either

L1. the image of γ lies on the y_0-axis, in which case

$$z = \lim_{t\to 0} \Psi(0, t) = \lim_{t\to 0}(0 : t^{cp} : 0 : t^{\beta_1} : t^{\beta_2}) = (0 : 0 : 0 : 0 : 1) =: P,$$

since $cp = p(r + s)(r + s_1) > (p - q)(r + s)r = \beta_1 > (p - q)(r + s_1)r = \beta_2$, or

L2. γ has a parametrization of the form $(t, \phi(t))$, where $\phi(t)$ is a Puiseux series in t.

Let $z \in E \setminus \{P\}$. Let $\phi(t) := \sum_{k=1}^{\infty} b_k t^{r_k}$ be as in assertion L2. Then

$$\Psi(\gamma(t)) = \left(t^{cpr} : b_1^{cp} t^{cpr_1} + \text{h.o.t.} : t^{cr(p-q)} : b_1^{\beta_1} t^{\beta_1 r_1} + \text{h.o.t.} : (\sum_{k=1}^{\infty} b_k t^{r_k} - \sum_{k=1}^{l} a_k t^{r+s_k})^{\beta_2} \right),$$

where h.o.t. is a shortcut for 'higher order terms' (in t). Note that $cr(p - q) < cpr$. Moreover, as proved in assertion L1 above, $cp > \beta_1$ and therefore, $cpr_1 > \beta_1 r_1$ (since $r_1 > 0$). Consequently,

$$z = \lim_{t\to 0} \Psi(\gamma(t)) = \lim_{t\to 0} \left(0 : 0 : t^{cr(p-q)} : b_1^{\beta_1} t^{\beta_1 r_1} : (\sum_{k=1}^{\infty} b_k t^{r_k} - \sum_{k=1}^{l} a_k t^{r+s_k})^{\beta_2} \right).$$

Let $\tilde{\phi}(t) := \sum_{k=1}^{\infty} b_k t^{r_k} - \sum_{k=1}^{l} a_k t^{r+s_k}$. We now show that $r_1 = r + s_1$ and $b_1 = a_1$. Indeed, if $r_1 < r + s_1$, then $\operatorname{ord}(\tilde{\phi}(t)) = r_1$. Moreover, $cr(p-q) = r(r+s)(r+s_1)(p-q) > r(r+s)r_1(p-q) = \beta_1 r_1 > \beta_2 r_1 = \beta_2 \operatorname{ord} \tilde{\phi}(t)$ (the last inequality is a consequence of the fact that $\beta_1 > \beta_2$, which we have shown in assertion L1 above). It follows that in this case $z = \lim_{t\to 0} \Psi(\gamma(t)) = P$, which is a contradiction to our choice of z. Similarly, if $r_1 > r + s_1$, or if $r_1 = r + s_1$ and $b_1 \neq a_1$, then $\operatorname{ord}(\tilde{\phi}(t)) = r + s_1$ and therefore $\beta_1 r_1 = r(r+s)(p-q)r_1 \geq r(r+s)(p-q)(r+s_1) = cr(p-q) > r(p-q)(r+s_1)(r+s_1) = \beta_2 \operatorname{ord} \tilde{\phi}(t)$ (here the last inequality is the consequence of the fact that $s > s_l \geq s_1$). Consequently, in this case as well, $z = \lim_{t\to 0} \Psi(\gamma(t)) = P$ which is a contradiction as noted above.

Since $r_1 = r + s_1$, it follows that $cr(p-q) = \beta_1 r_1 = r(p-q)(r+s)(r+s_1) = \beta_2(r+s)$. Therefore $\operatorname{ord} \tilde{\phi}(t) \geq r + s$, for otherwise we will again have that $z = P$. It follows that

$$\phi(t) = \sum_{k=1}^{l} a_k t^{r+s_k} + b t^{r+s} + \text{h.o.t.} \tag{43}$$

for some $b \in \mathbb{C}$. Consequently,

$$z = \lim_{t\to 0} \Big(0 : 0 : t^{cr(p-q)} : a_1^{\beta_1} t^{cr(p-q)} : b^{\beta_2} t^{cr(p-q)} + \text{h.o.t.}\Big) = \Big(0 : 0 : 1 : a_1^{\beta_1} : b^{\beta_2}\Big) \tag{44}$$

It follows that

$$E = \{(0 : 0 : 1 : a_1^{\beta_1} : \tilde{b}) : \tilde{b} \in \mathbb{C}\} \cup \{P\}. \tag{45}$$

Then $E \cong \mathbb{P}^1$ and therefore E is irreducible, which proves assertion 1 of the claim.

It follows from (43) that the Puiseux expansion for y_0 (in terms of w_0) associated with γ is

$$y_0 = \sum_{k=1}^{l} a_k w_0^{r+s_k} + b w_0^{r+s} + \text{h.o.t.}$$

In terms of y and x the above expression becomes

$$\begin{aligned}
\frac{y}{x^r} &= \sum_{k=1}^{l} a_k x^{-(r+s_k)} + b x^{-(r+s)} + \text{l.o.t.} \\
\Rightarrow y &= \sum_{k=1}^{l} a_k x^{-s_k} + b x^{-s} + \text{l.o.t.} \\
\Rightarrow y &= \psi_b(x) + \text{l.o.t.}
\end{aligned} \tag{46}$$

Identities (44), (45) and (46) prove assertions 2, 4 and 5 of the claim. It remains to prove that P is the intersection of E and the closure of $\Psi(X_\infty^{\delta_1})$ in $\bar{\Gamma}$. But we have shown in L1 that P is the point of intersection of E and the closure in $\bar{\Gamma}$ of the image of y_0-axis under Ψ. Since y_0-axis is precisely $X_\infty^{\delta_1}$, the proof of the claim is complete. ☐

We now show that E remains irreducible in the normalization of $\bar{\Gamma}$. More precisely, let Z be the normalization of $\bar{\Gamma}$, with the normalization map $\pi : Z \to \bar{\Gamma}$. Then

Claim 5.7.2. *$\tilde{E} := \pi^{-1}(E)$ is an irreducible curve. The discrete valuation on A associated to $\tilde{E}$ is $-\delta_2/p$.*

Proof. Let E' be an irreducible component of $\tilde{E}$ and ν' be the discrete valuation on A corresponding to E'. Let

$$h := x^s(y - \psi(x)) \in \mathbb{C}[x, y].$$

At first we show that for every $a \in \mathbb{C}$, $\nu'(h - a) = 0$. Indeed, if $\nu'(h - a) > 0$ for some $a \in \mathbb{C}$, then pick a Zariski open subset U' of Z such that $U' \cap E' \neq \emptyset$ and $h - a$ is regular on U'. Then h is also a regular function on U and $h(z) = a$ for all $z \in E' \cap U'$. Now, assertion 2 of claim 5.7.1 implies that there is $z \in E' \cap U'$ such that $\pi(z) = \lim_{t\to\infty} \gamma_b(t)$ for some $b \in \mathbb{C}$, $b \neq a$, where $\gamma_b(t) := (t, \psi(t) + bt^{-s})$ for all $t \in \mathbb{C} \setminus \{0\}$. Pick a curve $\tilde{\gamma}$ on $\pi^{-1}(\mathbb{C}^2)$ such that $\lim_{t\to\infty} \tilde{\gamma}(t) = z$. Assertion 5 of claim 5.7.1 implies that in (x, y)-coordinates $\pi(\tilde{\gamma}(t))$ has the following form:

$$\pi(\tilde{\gamma}(t)) = (t, \psi(t) + bt^{-s} + \text{l.o.t.}).$$

It follows that $h(z) = \lim_{t\to\infty} h(\tilde{\gamma}(t)) = \lim_{t\to\infty} h(\pi(\tilde{\gamma}(t))) = b$, which is a contradiction, since $b \neq a$. Therefore $\nu'(h - a) \leq 0$ for all $a \in \mathbb{C}$.

Similarly, if $\nu'(h - a)$ is negative for some $a \in \mathbb{C}$, then $\nu'(h)$ is also negative and we may pick a Zariski open subset U' of Z such that $U' \cap E' \neq \emptyset$ and h^{-1} is regular on U' and $h^{-1}(z) = 0$ for all $z \in E' \cap U'$. Choose $z \in E' \cap U'$ such that $\pi(z) = \lim_{t\to\infty} \gamma_b(t)$ for some $b \in \mathbb{C}$, $b \neq 0$. Then exactly as in the previous paragraph, we may show that $h^{-1}(z) = b^{-1} \neq 0$, which is a contradiction. It follows that for all $a \in \mathbb{C}$, $\nu'(h - a) = 0$, as required.

It follows that for all $a \in \mathbb{C}$,

$$\nu'(y - \psi(x) - ax^{-s}) = -s\nu'(x). \tag{47}$$

In particular, setting $a = 0$, we see that

$$\nu'(y - \psi(x)) = -s\nu'(x). \tag{48}$$

Recall that the projection from $\bar{\Gamma}$ to $\bar{X}$ maps E to $O \in \bar{X}$ and $w_0 := 1/x$ is regular at O. It follows that $1/x$ is regular at every point of $\tilde{E}$ and therefore $\nu'(x) \leq 0$. We claim that $\nu'(x) < 0$. Indeed, if $\nu'(x) = 0$, then (48) implies that $\nu'(y) \geq 0$, which implies that $\nu'(f) \geq 0$ for all $f \in \mathbb{C}[x, y]$. But this is impossible since $\tilde{E}$ is a curve at *infinity* with respect to $\mathbb{C}^2$ (see, e.g., [2, Proposition 1]). Therefore $\nu'(x) < 0$.

Let $\delta' := -\nu'$. Then δ' satisfies the hypothesis of corollary 2.7 and therefore there is a finite Puiseux series $\tilde{\psi}$ in x and a rational number $\tilde{r} < \operatorname{ord}_x(\tilde{\psi})$ such that for all $f \in \mathbb{C}[x, y]$,

$$\delta'(f) = \delta'(x) \deg_x(f(x, \tilde{\psi}(x) + \xi x^{\tilde{r}})). \tag{49}$$

Identities (48) and (47) then imply that

$$\deg_x(\tilde{\psi}(x) + \xi x^{\tilde{r}} - \psi(x)) = -s, \text{ and} \tag{50}$$

$$\deg_x(\tilde{\psi}(x) + \xi x^{\tilde{r}} - \psi(x) - ax^{-s}) = -s \text{ for all } a \in \mathbb{C}. \tag{51}$$

Since $\psi(x) = \sum_{k=1}^{l} a_k x^{-s_k}$ with $s_1 < s_2 < \cdots < s_l < s$, identities (50) and (51) imply that $\tilde{\psi} = \psi$ and $\tilde{r} = s$. It follows that δ' and δ_2 are proportional (remark 2.10). Also note that for each $f \in \mathbb{C}[x, y]$, $\delta'(f)$ is a multiple of $\delta'(x)$ (this follows from (49) since $\psi(x) \in \mathbb{C}[x, x^{-1}, y]$ and s is an integer). It follows that $\delta'(x) = 1$ and consequently

$$\delta' = \frac{\delta'(x)}{\delta_2(x)}\delta_2 = \frac{1}{p}\delta_2.$$

Therefore we have shown that every irreducible component of $\tilde{E}$ corresponds to the same discrete valuation on $\mathbb{C}(x, y)$. Since the discrete valuation associated to a curve uniquely distinguishes the curve from other curves on a surface, it follows that $\tilde{E}$ is irreducible and the discrete valuation associated to $\tilde{E}$ is precisely $-\delta_2/p$. ☐

Claim 5.7.2 implies that $Z_\infty := Z \setminus \mathbb{C}^2$ has precisely two components - one corresponding to each δ_i, $1 \leq i \leq 2$. Consequently, if $\tilde{\delta}$ is a subdegree on $\mathbb{C}[x, y]$ such that the corresponding completion $X^{\tilde{\delta}}$ of $\mathbb{C}^2$ is isomrphic to Z, then $\tilde{\delta} = \max\{r_1\delta_1, r_2\delta_2\}$ for some $r_1, r_2 \in \mathbb{Z}_{>0}$. The special case of the theorem will be proved if we show that there is a projective embedding of Z for which $r_1 = r_2$.

Recall that Z is the normalization of $\bar{\Gamma}$, where $\bar{\Gamma}$ is the closure of

$$\Gamma := \{(z, \Psi(z)) : z \in \mathbb{C}^2\} \subseteq X^{\delta_1} \times \mathbb{P}^4.$$

Therefore, choosing different projective embeddings of X^{δ_1}, we get different projective embeddings of $X^{\delta_1} \times \mathbb{P}^4$, and these in turn induces different projective embeddings of $\bar{\Gamma}$. In particular, let ϕ_d be the dr-uple embedding of X^{δ_1}, i.e.

$$\phi_d(W : X : Y) := (\cdots : W^\omega X^\alpha Y^\beta : \cdots)$$

where the monomials which appear as homogeneous coordinates in the right hand side of the above expression are precisely $S := \{W^\omega X^\alpha Y^\beta : \omega + \alpha + r\beta = dr\}$. Then composing $\phi_d \times \bar{\Gamma}$ is isomorphic to the closure in $\mathbb{P}^{|S|-1} \times \mathbb{P}^4$ of

$$\begin{aligned}\tilde{\phi}_d(x, y) &= \sigma_d(\phi_d(x, y), \Psi(x, y))\\ &= \sigma_d((1 : y : \cdots : y^d : \cdots : x^{dr}), (1 : g_1 : g_2 : g_3 : g_4))\\ &= (1 : g_1 : g_2 : g_3 : g_4 : \cdots : x^{dr} : x^{dr}g_1 : x^{dr}g_2 : x^{dr}g_3 : x^{dr}g_4)\end{aligned}$$

Let $U_\alpha := \mathbb{P}^{|S|-1} \setminus V(z_\alpha)$ and $V_j := \mathbb{P}^6 \setminus V(u_j)$. Recall that in the proof of claim 5.7.1 we computed the coordinates of the point P of intersection of the components of $\bar{\Gamma} \setminus \mathbb{C}^2$. In particular, $P \in U_{(0,dr,0)} \times V_4$. Let $\delta^{(d)}$ be a *normalization* of the degree-like function on $\mathbb{C}[x, y]$ corresponding to the embedding $\tilde{\phi}_d : \mathbb{C}^2 \hookrightarrow \bar{\Gamma}$. Then there exists a positive integer k_d such that the k_d-uple divisor at infinity on $X^{\delta^{(d)}}$ is generated by $1/(h_\alpha g_i)$ on $\pi^{-1}(\bar{\Gamma} \cap (U_\alpha \times V_j))$. This implies that

$$D^{\delta^{(d)}}_{k_d,\infty} = -\nu_1(g_4 x^{dr})[X_{\infty,1}] + -\nu_2(g_4 x^{dr})[X_{\infty,2}]$$

Since $g_4 = f_2$, it follows from (40) that $\nu_1(g_4) = -pr(r + s)(r + s_1)$ and $\nu_2(g_4) = -qr(r + s)(r + s_1)$. Recall that $\nu_1(x) = \nu_2(y) = -1$. It follows that

$$D^{\delta^{(d)}}_{k_d,\infty} = r(d + p(r + s)(r + s_1))[X_{\infty,1}] + r(d + q(r + s)(r + s_1))[X_{\infty,2}] \tag{52}$$

Now pick two positive integers p' and q' such that $p' > q'$. Note that replacing p and q respectively by p' and q' gives rise to a variety isomorphic to Z [3, Proposition 6.5]. Replacing p and q by suitable p' and q' and choosing a suitable d in (52) we may ensure that

$$D^{\delta^{(d)}}_{k_d,\infty} = mp[X_{\infty,1}] + mq[X_{\infty,2}]$$

for some positive integer m. But then $\delta^{(d)} = \max\{k\delta_1, k\delta_2\} = k\delta$ for some $k \geq 1$. Hence $k\delta$ and therefore δ is finitely generated. This completes the proof of the theorem in the special case. □

References

[1] Charles Favre and Mattias Jonsson. *The valuative tree*, volume 1853 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2004.

[2] Jacob Eli Goodman. Affine open subsets of algebraic varieties and ample divisors. *Ann. of Math. (2)*, 89:160–183, 1969.

[3] Pinaki Mondal. Projective completions of affine varieties determined via 'degree-like' functions. arXiv:1012.0835 [Math.AG], 2010.

[4] Pinaki Mondal. Primitive normal completions of the affine plane i. 2011.

[5] Stefan Schröer. On contractible curves on normal surfaces. *J. Reine Angew. Math.*, 524:1–15, 2000.

[6] Oscar Zariski and Pierre Samuel. *Commutative algebra. Vol. II.* Springer-Verlag, New York, 1975. Reprint of the 1960 edition, Graduate Texts in Mathematics, Vol. 29.